

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 223 219 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

17.07.2002 Bulletin 2002/29

(21) Application number: 02004252.9

(22) Date of filing: 30.10.1998

(51) Int Cl.7: **C12N 15/31**, C07K 14/39,
C12N 15/81, C12N 1/19,
C12P 21/02
// (C12N15/31, C12R1:645),
(C12N1/19, C12R1:645)

(84) Designated Contracting States:
DE FR GB

(30) Priority: 31.10.1997 JP 31460897

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
98950470.9 / 0 972 834

(71) Applicant: **ASAHI GLASS COMPANY LTD.**
Tokyo 100-8405 (JP)

(72) Inventors:

- Tohda, Hideki, Asahi Glass Company Ltd.
Yokohama-shi, Kanagawa 221-0863 (JP)

- Hama, Yuko, Asahi Glass Company Ltd.
Yokohama-shi, Kanagawa 221-0863 (JP)
- Kumagai, Hiromichi, Asahi Glass Company Ltd.
Yokohama-shi, Kanagawa 221-0863 (JP)

(74) Representative: **Müller-Boré & Partner**
Patentanwälte
Grafinger Strasse 2
81671 München (DE)

Remarks:

This application was filed on 26 - 02 - 2002 as a
divisional application to the application mentioned
under INID code 62.

(54) **Inducible promoter and secretion signal for use in *Schizosaccharomyces pombe*, expression vector containing them and their use**

(57) The present invention relates to an inducible promoter gene for use in the fission yeast *Schizosaccharomyces pombe*, an inducible expression vector containing the inducible promoter gene, *Schizosaccharomyces pombe* transformed with the inducible expression vector and a process for producing a heterologous protein by using the *Schizosaccharomyces pombe* transformant.

The present invention also relates to a secretion signal gene for use in *Schizosaccharomyces pombe*, an expression vector containing the secretion signal, *Schizosaccharomyces pombe* transformed with the expression vector and a process for producing a heterologous protein by using the *Schizosaccharomyces pombe* transformant. These inducible promoter gene and secretion signal gene are originated from the *S. pombe* gene involved in expression of the invertase precursor.

The inducible promoter gene is originated from the region responsible for the catabolite repression of the

invertase precursor. When a *Schizosaccharomyces pombe* transformant carrying the inducible expression vector is incubated, the timing of the production of the desired heterologous protein can be controlled by regulating the expression of the heterologous protein structural gene depending on the presence or absence of a specific nutrient (especially glucose).

The secretion signal gene encodes the secretion signal for the invertase precursor. When a *Schizosaccharomyces pombe* transformant carrying an expression vector containing the secretion signal gene is incubated, a heterologous protein fused with the secretion signal is synthesized as a precursory protein in the cell, and after intracellular separation of the secretion signal, the heterologous protein is secreted from the cell. Therefore, it is possible to obtain the desired protein from the cell culture, and the desired heterologous protein can be purified easily.

EP 1 223 219 A2

Description

[0001] The present invention relates to an inducible promoter gene and secretion signal gene for use in the fission yeast Schizosaccharomyces pombe (hereinafter referred to as S. pombe), an expression vector containing them and a process for producing a protein using them. In particular, it relates to a process for producing a desired protein wherein the S. pombe invertase promoter is used to make it possible to control the timing of the protein production by the presence or absence of a specific nutrient through regulated gene expression, and a process for secretory production of a desired protein by using the secretion signal gene for the S. pombe invertase precursor.

[0002] S. pombe, despite being a eukaryote, has been studied extensively for its high versatility in genetics, molecular biology and cellular biology as a unicellular organism (Nasim A. et al. eds., Molecular biology of the fission yeast, Academic Press, 1989). In its cultures, monosaccharides such as glucose and fructose are used as the main carbon sources. It is known that in a culture medium lacking these monosaccharides, expression of invertase, the enzyme that degrades sucrose into glucose and fructose, is induced to secure the carbon source necessary for its growth (Moreno S. et al., Arch Microbial. 142, 370, 1985).

[0003] S. pombe invertase is located on the cell surface and is a high-molecular weight glycoprotein with a molecular weight of about 205000, 67% of which is attributed to sugar chains composed of equimolar amounts of mannose and galactose residues. Molecular weight and amino acid studies of the protein moiety of the pure enzyme and experiments using antibodies have shown high similarity between S. pombe invertase and the invertase from the budding yeast Saccharomyces cerevisiae from the viewpoint of protein chemistry (Moreno S. et al., Biochem. J. 267, 697, 1990). It is also known that a drop in glucose concentration de-represses synthesis of invertase (Mitchinson J. et al., Cell Sci 5, 373, 1969).

[0004] Induced invertase synthesis (de-repression) is also observed in Saccharomyces cerevisiae. Previous detailed studies on genetic regulation of invertase expression, the biosynthetic pathway and the structure of the sugar chain moiety have shown that Saccharomyces cerevisiae invertase is encoded by six overlapping genes, SUC1 to SUC5 and SUC7, on one chromosome and that activation of at least one of these SUC genes leads to utilization of sucrose and raffinose (Hohmann S. et al., Curr Genet 11, 217, 1986).

[0005] In contrast, with respect to S. pombe, although purification of the invertase protein has been reported (Moreno S. et al. 1985), no invertase genes had been identified until the present inventors and coworkers recently reported two overlapping invertase genes *inv0+* and *inv1+* in S. pombe. Because *inv0+* is likely a pseudogene having an incomplete open reading frame, *inv1+* is the only one gene encoding S. pombe invertase, which is supposed to confer the ability to grow on sucrose even in the absence of other carbon sources ("Kobogaku" edited by Yositaka Hashitani, Iwanami Shoten, 1967).

[0006] Analysis of the promoter region of the isolated gene suggested that a specific sequence between the 1st and 62nd base pairs is involved in catabolite repression.

[0007] In Saccharomyces cerevisiae, the SUC2 gene is transcribed into two messenger RNAs (mRNAs) from different transcription initiation sites. The shorter one is a constitutive mRNA encoding the intracellular invertase, while the longer one is a mRNA encoding the catabolite-repressible secretory invertase with a de-repression ratio of not less than 200 (Carlson M. et al., Mol. Cell. Biol. 3, 439, 1983). Analysis of the promoter region for the longer mRNA suggested that the transcription initiation factor binds to a specific repeated sequence between positions -650 and -418 (Salokin L et al., Mol. Cell. Biol. 6, 2314, 1986). The region between positions -418 and -140 has been shown to be necessary for glucose repression.

[0008] These regions in the SUC2 gene showed no significant homology with the *inv1+* upstream region between positions 1 and 2809. However, multiple copies of a so-called 7-bp motif with the sequence (A/C) (A/G)GAAAT, which is repeated at five sites in the region indispensable for glucose derepression, have been found in the *inv1+* upstream region. Further, while palindrome stem-loops have been identified at almost the same positions in the upstream regions of glucose-repressible genes (SUC, MAL and GAL), palindrome sequences have also been found in the upstream region of the *inv1+* gene from S. pombe. These sequences are anticipated to play an important role in glucose repression in S. pombe.

[0009] The yeast S. pombe is phylogenetically different from Saccharomyces cerevisiae. It is quite different from other yeasts in the chromosome structure and various mechanisms for genome replication, RNA splicing, transcription and posttranslational modification, and rather resembles animal cells in some of these aspects. For this reason, S. pombe is widely used as a eukaryotic model (Giga-Hama and Kumagai, eds., Foreign gene expression in fission yeast Schizosaccharomyces pombe, Springer-Verlag, 1997).

[0010] S. pombe is also widely used as a host for expression of heterologous protein genes and known to be suited especially for expression of genes from animals including human (JP-A-5-15380 and JP-A-7-163373). For its advanced membrane structures including the Golgi body and the endoplasmic reticulum, S. pombe is also used for expression of membrane proteins and shows high level expression. For S. pombe, constitutive expression vectors (pEVP11, pART1 and pTL2M) and an inducible expression vector using the promoter region of the *nmt1+* gene (pREP1) are usually

used as expression vectors. No S. pombe expression vectors of the GAL type or the SUC type have been known though these types of vectors are commonly used for Saccharomyces cerevisiae.

[0011] The expression of the SUC2 gene from Saccharomyces cerevisiae in S. pombe has been shown to be constitutive, not catabolite repressible, though the expression product contains galactose residues conferred by the host (Zarate, V. et al., J Applied Bacteriology, 80, 45, 1996), suggesting differences between S. pombe and Saccharomyces cerevisiae in the mechanism for catabolite repression of invertase. The differences are of great significance because the promoter from Saccharomyces cerevisiae usually used by those skilled in the art for construction of inducible expression vectors of the invertase type (the SUC2 type) is not applicable to S. pombe vectors. Therefore, development of S. pombe vectors of this type has been long delayed.

[0012] On the other hand, the present inventors constructed an expression vector using the secretion signal gene encoding the secretion signal in the precursor of a S. pombe mating pheromone (WO96/23890). However, this secretion signal gene is not an all-purpose secretion signal gene, and other secretion signal genes that function in S. pombe are desired for production of some types of protein.

[0013] As a result of their extensive research with a view to solving the above problems, the present inventors have accomplished the present invention by preparing a new clone of the S. pombe invertase gene and constructing an inducible expression vector. They have also found that the N-terminal 22 amino acid sequence in the amino acid of the invertase precursor functions as a secretion signal. On the basis of these findings, they have constructed an expression vector using the secretion signal gene and established secretory production of desired proteins.

[0014] The present invention relates to a region in the invertase gene from S. pombe, which is involved in catabolite repression, an inducible expression vector using the region and a system using it for heterologous gene expression and provides:

a DNA in an invertase gene from Schizosaccharomyces pombe, which is located in a region involved in catabolite repression,

a DNA having the base sequence of bases 1 to 2809 in SEQ ID NO: 1 in the Sequence Listing,

a recombinant vector containing the sequence of the DNA, a multicloning vector containing the sequence of the DNA and a multicloning site,

a multicloning vector having the structure shown in Figure 9,

an expression vector for transformation of Schizosaccharomyces pombe containing the sequence of the DNA and

a heterologous protein structural gene,

a transformant from Schizosaccharomyces pombe containing the expression vector, and

a process for producing a protein which comprises incubating the transformant and recovering an expressed heterologous protein.

[0015] Firstly, the present inventors cloned and sequenced a S. pombe invertase gene, which had not been genetically identified. Then, they demonstrated by gene disruption analysis that the invertase gene is responsible for the overall invertase activity. Further, they identified the region involved in catabolite repression and constructed an inducible expression vector using the region. They actually constructed a recombinant vector carrying the gene of a green fluorescent protein, transformed S. pombe with the vector and confirmed the expression of the protein by assay of invertase activity and immunological analysis. They also demonstrated repression of the heterologous gene expression in the presence of glucose in the culture medium and derepression by exhaustion of glucose.

[0016] The present inventors used the following procedure to identify and characterize the gene of the S. pombe invertase precursor:

(1) PCR using a cDNA library from S. pombe as a template and primers based on conserved amino acid sequences in invertase genes from many other organisms;

(2) screening of a genomic library from S. pombe by plaque hybridization using the PCR product as a probe for positive clones;

(3) confirmation of the positive clones by restriction digestion followed by electrophoresis;

(4) Southern hybridization analysis and total sequencing of a fragment with a specific length in the positive clones;

(5) gene disruption analysis of invertase activity;

(6) investigation of the optimum pH for expression of the invertase gene from S. pombe and the effects of the glucose concentration on glucose repression and derepression; and

(7) identification of a region indispensable for glucose repression through subcloning of the related upstream region.

[0017] Also, the present inventors constructed a S. pombe invertase inducible expression vector by the following procedure and actually demonstrated inducible expression of a green fluorescent protein:

- (1) construction of an inducible multicloning expression vector pRI0M containing an invertase promoter by modifying a S. pombe multicloning vector, pTL2M (JP-A-7-163373);
- (2) construction of an inducible expression vector pRI0EGFP for expression of a green fluorescent protein variant from the inducible multicloning vector, pRI0M;
- (3) transformation of a wild-type S. pombe strain with the inducible expression vector, pRI0EGFP, for expression of the green fluorescent protein variant;
- (4) demonstration of the expression of the green fluorescent protein variant by activity (fluorescence) analysis and SDS-PAGE-western blotting; and
- (5) establishment of the conditions for the inducible expression on the basis of the dependence of the expression level on the glucose concentration in the culture medium.

[0018] SEQ ID NO: 1 in the Sequence Listing is the base sequence of the gene of the invertase precursor, which contains a region involved in catabolite repression. The region involved in catabolite repression is the DNA sequence between positions 1 to 2809 or within the DNA sequence. In the DNA sequence between positions 1 and 2809, the region extending from position 1 to position 620 and the region extending from position 1610 to position 2610 are especially important, as is evident from the results of the analysis in Example 6 shown in Figure 8 (position 2810 in SEQ ID NO: 1 corresponds to position 1 in Figure 8). This means that the inducible promoter in the present invention is not restricted to a DNA having the base sequence from position 1 to position 2809 so long as it contains these genes involved in catabolite repression and functions as an inducible promoter. Still, a DNA having a base sequence from position 1 to position 2809 is preferred as an inducible promoter because it actually functions in S. pombe.

[0019] The above-mentioned DNA which contains genes involved in catabolite repression and function as an inducible promoter, preferably having the base sequence from position 1 to position 2809, is hereinafter referred to as the inducible promoter gene. The inducible promoter gene can be integrated with a vector for construction of recombinant vectors such as multicloning vectors and expression vectors. A multicloning vector is a vector having a multicloning site and provides an expression vector through introduction of a desired structural gene into the multicloning site. An expression vector is a vector containing a structural gene and used for expression of a structural gene encoding a heterologous protein. A "heterologous" protein is a protein which is not inherent in the host. For example, when the host is S. pombe, a heterologous protein is a protein which is not inherent in S. pombe (such as a human protein).

[0020] In the expression vector, the inducible promoter gene is located upstream from the heterologous protein structural gene and regulates expression of the structural gene. The inducible promoter gene in the expression vector regulates the expression of the heterologous protein structural gene downstream, like the inducible promoter gene, located upstream in the base sequence represented by SEQ ID NO: 1 regulates the expression of the structural gene of the invertase precursor. In the multicloning vector, the inducible promoter gene is located upstream from the multicloning site into which a heterologous protein structural gene is to be introduced.

[0021] One example of the multicloning vector of the present invention is the multicloning vector pRI0M constructed in Example 7 and has the structure shown in Figure 9. The entire base sequence of pRI0M is SEQ ID NO: 2. Inv1-P is the above-mentioned inducible promoter gene, and MCS is the multicloning site. One example of the expression vector of the present invention is the inducible expression vector pRI0EGFP for expression of a green fluorescent protein variant constructed in Example 8 by introducing the structural gene (EGFP-ORF) of a green fluorescent protein variant and has the structure shown in Figure 10. The entire base sequence of the expression vector pRI0EGFP is SEQ ID NO: 13.

[0022] The most suitable cell (host) to transform with the expression vector of the present invention is S. pombe because the inducible promoter gene in the present invention is an inducible promoter gene from S. pombe.

[0023] Under catabolite repressing conditions (for example, in a culture medium containing a high level of glucose), S. pombe transformed with the expression vector of the present invention grows with no (or low) expression of the heterologous protein. Growth at this stage without the burden of heterologous protein expression is more efficient than growth under the burden. Subsequent incubation under catabolite derepressing conditions (for example, in a culture medium containing no or a low level of glucose) invites the increased number of S. pombe cells to high level expression of the heterologous protein, though growth of S. pombe is less efficient than under catabolite repressing conditions. Thus, controlled transition between growth of S. pombe and heterologous protein expression through catabolite repression allows more efficient production of a heterologous protein.

[0024] Catabolite repression can be controlled not only in an active way as described above but also in a passive way. For example, when a S. pombe transformant is incubated in a culture medium containing a given amount of glucose, the S. pombe grows under catabolite repressing conditions containing a high level of glucose in the initial stage, but later on production of a heterologous protein predominates due to catabolite derepression as glucose is exhausted. This way, more efficient heterologous protein production of than ever is possible without active control of the glucose level.

[0025] In addition to the above-mentioned total sequencing of the invertase precursor gene from S. pombe, the

present inventors determined the complete amino acid sequence of the invertase precursor (the amino acid sequence in SEQ ID NO: 1). Then, they have found that the first 22 amino acid peptide in the amino acid sequence of the invertase precursor (Met Phe Leu Lys Tyr Ile Leu Ala Ser Gly Ile Cys Leu Val Ser Leu Leu Ser Ser Thr Asn Ala) acts as a secretion signal. Hereinafter, the peptide is referred to as the secretion signal.

[0026] It is expected that a desired heterologous protein produced in transformed *S. pombe* cells as a protein fusion having the secretion signal at the N-terminal is secreted from the cells after intracellular processing which splits the protein fusion into the secretion signal and the heterologous protein. The present inventors constructed an expression vector carrying a heterologous protein structural gene (specifically, human interleukin 6-a'c1 variant) fused with a DNA encoding the secretion signal (namely, a structural gene of a protein fusion as mentioned above) and demonstrated secretion of the heterologous protein from *S. pombe* cells transformed with the expression vector.

[0027] The present invention provides the secretion signal, a DNA encoding the secretion signal (hereinafter referred to as a secretion signal gene), a recombinant vector carrying the secretion signal gene, a multicloning vector carrying the secretion signal gene, an expression vector carrying the secretion signal gene and a heterologous protein structural gene for transformation of *S. pombe*, a *S. pombe* transformant carrying the expression vector and a process for producing a protein which comprises incubating the transformant and recovering the expressed heterologous protein.

[0028] The secretion signal gene is not restricted to the 66-bp sequence extending from position 2810 to position 2875 in SEQ ID NO: 1 and may be a DNA having a different base sequence encoding the amino acid sequence of the secretion signal. In the expression vector, the secretion signal and the heterologous protein structural gene is preferably linked directly. But they may be linked via another DNA sequence, for example, extending from position 2876 in SEQ ID NO: 1. In this case, the protein product has extra amino acid residues at the N-terminal of the heterologous protein but can be converted into the desired heterologous protein by trimming off the N-terminal extra amino acid residues. However, the disadvantage from the presence of these extra amino acid residues usually becomes more serious for the desired protein as the number of extra amino acid residues increases. Therefore, as the intervening DNA between the secretion signal gene and the heterologous protein structural gene, a short DNA encoding at most 10 amino acid residues is preferable. Particular preferably, the secretion signal gene and the heterologous protein structural gene are linked directly.

[0029] Construction of an expression vector carrying the secretion signal gene using a multicloning vector can be attained by inserting a heterologous protein structural gene fused with the secretion signal gene into the multicloning site of a multicloning vector or by inserting a heterologous protein structural gene into the multicloning site of a multicloning vector carrying the secretion signal gene. The latter method tends to restrict the structure of the multicloning site because the secretion signal gene is preferred to be located immediately in front of the multicloning site as described above. Therefore, the former method is preferred for construction of an expression vector. As a *S. pombe* multicloning vector, for example, pTL2M, which is disclosed in JP-A-163373, is preferable.

[0030] According to the present invention, an expression vector can be constructed by using both the inducible promoter gene and the secretion signal gene. For example, an expression vector which contains the DNA sequence of from position 1 to position 2875 in SEQ ID No: 1 and a heterologous protein structural gene introduced downstream of the DNA sequence can be constructed. Such an expression vector enables catabolite repressible secretory production of a heterologous protein by the host cell. A similar expression vector can be constructed by using a known secretion signal gene (such as the secretion signal gene disclosed in WO96/23890) instead of the above-mentioned secretion signal gene.

[0031] Figure 1 shows a comparison of (partial) amino acid sequences deduced from *inv1*⁺, the *Schwanniomyces occidentalis* invertase gene and the fission yeast *SUC2* gene. Figure 2 is the restriction map of the *inv1*⁺ gene. Figure 3 electrophoretically shows disruption of the *inv1*⁺ gene. Figure 4(a) is a photograph of colony gel overlay assay of invertase activity (for phenotype characterization) as a substitute for a drawing. Figure 4(b) is a schematic explanation of experimental design of the invertase activity assay shown in Figure 4(a). Figure 5(a) is a photograph of colony gel overlay assay of invertase activity (for phenotype characterization) as a substitute for a drawing. Figure 5(b) is a schematic explanation of experimental design of the invertase activity assay shown in Figure 5(a).

[0032] Figure 6 graphically shows the relation between the invertase activity and the glucose concentration. Figure 7 graphically shows the relation between the invertase activity and the glucose concentration. Figure 8 shows the results of analysis of invertase promoters. Figure 9 show the structure of an inducible expression vector pRIOM. Figure 10 shows the structure of an inducible expression vector pRI0EGFP for expression of a green fluorescent protein.

[0033] Figures 11(a) and 11(b) demonstrate the expression of a green fluorescent protein. Figures 12(a) and 12(b) graphically show the relation between the incubation time and the expression level of the green fluorescent protein. Figures 13(a) and 13(b) graphically show the relation between the incubation time and the expression level of the green fluorescent protein. Figures 14(a) and 14(b) show the relation between the incubation time and the expression level of the green fluorescent protein. Figure 15 is a SDS-PAGE pattern obtained in analysis of the expression of interleukin 6a'c1 variant. Figure 16 is the western blot pattern of the expressed interleukin variant.

[0034] Now, the present invention will be described in further detail with reference to specific Examples.

EXAMPLE 1

Isolation of *S. pombe* invertase gene

[0035] PCR using a *S. pombe* cDNA library as the template and primers designed on the basis of conserved sequences in invertase genes from other organisms shown in SEQ ID NOS: 3 to 5, gave amplification products of about 300bp and about 400bp. Each PCR product was purified by using EASY TRAP (Takara Shuzo Co., Ltd.) and sequenced after ligation into a vector by using pMOS Blue T vector kit (Amersham Pharmacia Biotech K.K.). The deduced amino acid sequence indicated that part of the 400-bp PCR product has significant homology with the SUC2 gene from

Saccharomyces cerevisiae.

[0036] Screening of a genomic library of *S. pombe* for the entire invertase gene by plaque hybridization using the 400-bp PCR product as a probe picked up 15 positive clones from about 8,000 plaques. Secondary screening of the positive clones by plaque hybridization left four positive clones. Treatment of small amounts of phage DNA extracts from the positive clones with various restriction enzymes gave identical cleavage patterns, and thus revealed that all the clones were identical.

[0037] The entire invertase gene was isolated by the following two-step procedure. A 3.0-kb HindIII fragment was isolated from the hybrid-forming clones and ligated into a plasmid pBluescript II SK- (Toyobo Co., Ltd.). Restriction mapping identified BamHI and Sall sites. Subcloning of the fragment using these restriction enzymes and subsequent sequencing using a deletion technique revealed that the HindIII fragment contains the complete ORF of the gene but contains only about 200 bp within the upstream region, which is supposedly involved in the gene expression. Therefore, separately, a 3.5-kb BamHI fragment from the hybrid-forming clones was further digested with HindIII to give a 2.6-kb fragment. Subcloning of the 2.6-kb fragment in plasmid pBluescript II SK-, and subsequent sequencing using a deletion technique revealed that the BamHI-HindIII fragment contained a sequence within the upstream region supposedly involved in the gene expression. The resulting complete 5.6-kb gene was designated as *inv1+*. The base sequence of *inv1+* and the amino acid sequence encoded by its ORF are shown in SEQ ID NO: 1. A plasmid carrying the complete gene was designated as pINV3000.

[0038] These results suggest that the *inv1+* product has 16 asparagine-linked glycosylation sites. Figure 1 shows a comparison of the amino acid sequence deduced from the base sequence of the *inv1+* gene from *S. pombe*, the amino acid sequence of *Schwanniomyces occidentalis* invertase and the amino acid sequence deduced from the base sequence of the SUC2 gene from *Saccharomyces cerevisiae*. Amino acids that are common to the three are marked with *. Figure 1 clearly shows the amino acid sequence deduced from the *inv1+* base sequence has significant homology with invertases from other origins such as *Schwanniomyces occidentalis* and *Saccharomyces cerevisiae*, which suggests the *inv1+* encodes invertase.

EXAMPLE 2

Disruption of the *inv1+* gene

[0039] The HindIII site in plasmid pBluescript II SK- having a *S. pombe* *ura4+* gene insert at the ClaI site was disrupted by HindIII digestion followed by blunting and self-ligation (self-cyclization). Double restriction digestion of the plasmid with XbaI and HincII gave a *ura4+* fragment. The plasmid pBluescript was integrated with the *ura4+* fragment after restriction digestion with SpeI and subsequent blunting and XbaI digestion, to provide a plasmid having BamHI sites on both sides of *ura4+*. The BamHI site in plasmid pBluescript II SK- was disrupted similarly by restriction digestion followed by blunting and self-ligation, and the 3.0-kb fragment containing the *inv1+* ORF was inserted at the HindIII site. BamHI digestion of the plasmid eliminated a 1.4-kb fragment (containing part of the *inv1+* ORF encoding the C-terminal of invertase) from the 3.0-kb insert, and a *ura4+* cassette having BamHI sites at both ends was inserted. HindIII digestion of the resulting plasmid gave a DNA fragment having *inv1+* neighboring regions at both ends (Figure 2). The restriction map of the *inv1+* gene is shown in Figure 2, wherein the open reading frame (ORF) is indicated by the arrow (*inv1+*ORF) and the *ura4+* replacement from *Schizosaccharomyces pombe* is boxed (*ura4+*). The disruption mutant strain had an *inv1+* fragment carrying the *S. pombe* *ura4+* gene instead of the 1.4-kb *inv1+* BamHI-BamHI fragment partly containing the ORF. The *inv1+* fragment was used to transform a wild-type *S. pombe* strain, TP4-1D [*h⁻, leu1, ura4, ade6-M216, his2*, obtained from Dr. Takashi Toda (Imperial Cancer Research Foundation)], and viable colonies on a uracil-free culture medium were collected. Overlay assay of invertase activity revealed that 7 out of 28 strains, namely 25% of the *ura4+* colonies, lacked invertase activity.

[0040] Further, to verify the chromosomal *inv1+* gene disruption, genomic DNA from a strain lacking invertase activity was analyzed after double restriction digestion with HindIII and Sall by Southern hybridization using the *inv1+* HindIII-Sall fragment (2 kb) as the probe. The 3-kb hybridized fragment, which was not digested with Sall, shown in Figure 3 demonstrates that part of the *inv1+* gene in the chromosomal DNA had been replaced with the *ura4+* gene in the

strain which lacked invertase activity.

[0041] Thus, the *inv1⁺* gene proved to be the only one invertase gene expressed in *S. pombe*.

EXAMPLE 3

Restoration of invertase activity by the *inv1⁺* gene

[0042] The 3.0-kb HindIII fragment containing the entire *inv1⁺* ORF from the invertase gene was inserted into *S. pombe* vector pAU-SK (obtained from Dr. Chikashi Shimoda, Department of Science, Osaka City University), and the resulting recombinant vector was used to transform the invertase-defective strains (Example 2). The resulting transformants were streaked on YP sucrose plates (supplemented with 10 µg/ml antimycin A and 20 µg/ml bromocresol purple) for overlay assay of invertase activity. Further, the 2.6-kb *inv1⁺* BamHI-HindIII fragment containing the upstream promoter region and the 2.0-kb HindIII-Sall fragment from the invertase gene containing the ORF were ligated, and the resulting 4.6-kb BamHI-Sall fragment was inserted into pAU-SK. Transformation of the resulting recombinant vector into the invertase-defective strain was followed by similar overlay assay of invertase activity.

[0043] Both transformants (*inv1Δ*[pAU-SK::*inv1⁺*]) and the wild-type strain TP4-1D (WT) developed blue stains, which indicate invertase production, unlike the *inv1⁺* disruption mutant strain (*inv1Δ*). The addition of the upstream promoter region resulted in stronger stains, which suggest high-level invertase expression (Figure 4(a) and 4(b)). Figure 4(a) is a photograph showing the results of the gel overlay assays, and Figure 4(b) is a schematic explanation of the stained sections shown in Figure 4(a).

[0044] The invertase-defective strain were hardly viable on YP sucrose plates (supplemented with 10 µg/ml antimycin) whereas the wild-type strain and the transformants were recognizably viable after 5 days incubation at 30°C (Figure 5(a) and (b)). Figure 5(a) is a photograph showing the results of characterization by colony formation, and Figure 5(b) schematically explains the characterization shown in Figure 5(a).

[0045] These results demonstrate that the *inv1⁺* gene expression product is the invertase located on the cell surface.

EXAMPLE 4

Determination of glucose concentration for gene repression

[0046] For determination of the critical glucose concentration for catabolite repression of the invertase gene, the wild-type strain TP4-1D was incubated at 30°C in 5 ml of MM medium containing 2%, 4%, 8% and 16% glucose with shaking to the mid-logarithmic growth phase. Invertase assays were done by the method of Goldstein et al., and the post-incubational glucose concentrations in the medium were determined by the phenol-sulfate method (Figure 6). The hatched bars indicate invertase activity per cell (U/OD), and the empty bars indicate the residual glucose concentration. Judging from the graph, a glucose concentration of 8% is the optimum for glucose repressing incubation, because when the glucose concentration was 8%, the invertase activity was sufficiently repressed with little decrease of glucose.

EXAMPLE 5

Determination of glucose concentration for induced invertase production

[0047] For determination of the most effective glucose concentration for induced invertase production, the wild-type strain TP4-1D and a transformant [obtained by transforming the invertase-defective strain (Example 3) with a pAU-SK vector carrying the *inv1⁺* BamHI-Sall fragment] were preincubated in a medium containing 2% glucose to the mid-logarithmic growth phase and incubated in an MM medium containing 0%, 0.01%, 0.05%, 0.1%, 0.25%, 0.5%, 1.0% and 2% glucose with shaking at 27°C for 3 hours, and the invertase activity was assayed (Figure 7). Each run of assay was carried out at 30°C over 180 minutes. 1U of invertase converts 1 nmol of sucrose into glucose per 1 minute at 30°C, pH 4.0.

[0048] The optimum glucose concentration for induction was found to be 0.1% for the wild-type strain and 0.05% for the transformant. The invertase activity in the wild-type strain was 40 times higher under derepressing conditions than under repressing conditions. These results demonstrate catabolite repression in *S. pombe*.

EXAMPLE 6

Analysis of the inv1⁺ promoter region

5 [0049] Fragments obtained ligating *S. pombe* inv1⁺ upstream sequences extending from positions 1, 620, 1100, 1610 and 2610, respectively, and the inv1⁺ ORF were inserted into expression vector pAU-SK to obtain 5 plasmids for deletion studies. The plasmids containing the upstream sequences extending from positions 1, 1610 and 2610 carry the inv1⁺ BamHI-Sall, SacI-Sall and HindIII-Sall fragments, respectively. The plasmids containing the inv1⁺ upstream sequences extending from positions 620 and 1100 were constructed by site specific introduction of a SpeI site into

10 pAU-SK::inv1⁺ (BamHI-Sall) using primers shown in SEQ ID NOS: 6 and 7, respectively, followed by partial removal of the upstream region by SpeI treatment.

[0050] The plasmids thus obtained were used to transform the invertase-defective strain (Example 3). Invertase assays were done to determine the enzyme activity in each transformant (Fig. 8). The results suggest that the sequence between position 1 and position 620 is essential for glucose repression. The region between position 1620 and position

15 2610 was identified as essential for high-level glucose derepression of invertase.

EXAMPLE 7

Construction of inducible multicloning expression vector pRI0M carrying the invertase promoter

20 [0051] PCR amplification using the plasmid pINV3000 (Example 1) carrying the invertase gene from *S. pombe* as the template and oligo DNAs shown in SEQ ID NOS: 8 and 9 as the primers was performed to give a sequence which contains the promoter region for the invertase gene and has restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with SpeI (Takara Shuzo Co., Ltd.) and EcoRI (Takara Shuzo Co., Ltd.), the

25 sequence was subjected to agarose gel electrophoresis. Purification of the band of about 3000 bp by the glass beads method using EASY-TRAP gave an insert fragment.

[0052] The *S. pombe* multicloning vector pTL2M (JP-A-7-163373) was subjected to agarose gel electrophoresis after terminal double restriction digestion with SpeI and EcoI. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

30 [0053] The two fragments were ligated with a DNA ligation kit (Takara Shuzo Co., Ltd.) and transformed into *E. coli* strain DH (Toyobo Co., Ltd.). *E. coli* colonies were screened for the inducible expression vector pRI01M shown in Figure 9 and SEQ ID NO: 2 in the Sequence Listing through base sequencing and restriction mapping, and the inducible expression vector was recovered by the alkali-SDS method on a preparatory scale.

35 EXAMPLE 8

Construction of inducible expression vector pRI0EGFP for expression of green fluorescent protein

40 [0054] PCR amplification using the plasmid pINV3000 (Example 1) carrying the invertase gene from *S. pombe* as the template and oligo DNAs shown in SEQ ID NOS: 8 and 10 as the primers was performed to give a sequence which contains the promoter region for the invertase gene and has restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with SpeI and NheI (Takara Shuzo Co., Ltd.), the sequence was subjected to agarose gel electrophoresis. Purification of the band of about 3000 bp by the glass beads method using EASY-TRAP gave a fragment for use as the promoter insert.

45 [0055] PCR using the plasmid pEGFP carrying the jellyfish (*Aequorea victoria*) green fluorescent protein variant gene (Clontech) as the template and oligo DNAs shown in SEQ ID NOS: 11 and 12 in the Sequence Listing as the primers was performed to amplify the ORF in the green fluorescent protein variant gene. The PCR product was subjected to agarose gel electrophoresis after terminal double restriction digestion with NheI and HindIII. Purification of the band of about 700 bp by the glass beads method using EASY-TRAP gave a fragment for use as the ORF insert.

50 [0056] The *S. pombe* multicloning vector pTL2M was cleaved by double restriction digestion with SpeI and HindIII and then subjected to agarose gel electrophoresis. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

[0057] The three fragments were ligated with a DNA ligation kit and transformed into *E. coli* strain DH5. *E. coli* colonies were screened for the inducible expression vector pRI0EGFP for expression of the green fluorescent protein variant shown in Figure 10 and SEQ ID NO: 13 in the Sequence Listing through base sequencing and restriction mapping, and the inducible expression vector was recovered by the alkali-SDS method on a preparatory scale.

55

EXAMPLE 9

Preparation of *S. pombe* transformant ASP138

[0058] *S. pombe* wild-type strain ARC001 [leu1-32h⁻ (isogenic to ATCC38399)] was transformed with the inducible expression vector pRI0EGFP for expression of the green fluorescent protein variant and a transducing vector pAL7 as described by Okazaki et al. (Okazaki et al., "Nucleic Acids Res.", 18, 6485-6489, 1990).

[0059] 1 ml of a preincubated ARC001 culture in YPD medium was incubated in minimum medium MB+Leu with shaking at 30°C for 14 hours to a cell density of 3×10^7 per 1 ml, and the cells were collected, washed with water, suspended in 1 ml of 0.1M lithium acetate (pH 5.0) and incubated at 30°C for 60 minutes. A 100 µl portion of the suspension was mixed with 4 µg of the inducible expression vector pRI0EGFP and 0.5 µg of PstI-digested pAL7 in 15 µl TE and further with 290 µl of 50% PEG 4000 thoroughly and incubated at 30°C for 60 minutes, at 43°C for 15 minutes and at room temperature for 10 minutes, successively. After centrifugal removal of PEG, the cells were suspended in 1 ml of 1/2 YEL+Leu medium. After 10-fold dilution, 1 ml of the suspension was incubated at 32°C for 2 hours, and a 300 µl portion was spread on minimum medium agar MMA. After 3 days of incubation at 32 °C, about 300 independent colonies had developed on the plate.

[0060] 10 colonies of the transformant were inoculated in 2 ml of YEL medium containing 10 µg/ml antibiotic G418 (YEL10 medium) and incubated with shaking at 32°C. 2 days later, 6 clones were viable. In their subcultures, 4 clones were viable 3 days later. The putative desired transformant (ASP138 strain) was frozen in glycerol and stored for use in subsequent experiments.

EXAMPLE 10

Analysis of expression of green fluorescent protein variant

[0061] *S. pombe* transformant ASP138 (Example 9) was inoculated in YPD medium containing 100 µg/ml G418 (YPD100) and incubated at 32°C for 2 days. Green fluorescence was observed from each cell under a fluorescence microscope (excitation wavelength 490nm/emission wavelength 530nm). Green fluorescence emission was also observed upon ultraviolet irradiation from the centrifugally collected cells. Thus, expression of the desired green fluorescent protein in the active form was confirmed.

[0062] Strain ASP138 (Example 9) was incubated in 5 ml YPD100 at 32°C for 3 days, collected, washed and suspended in 50mM tris-HCl (pH 7.5), disrupted with glass beads in a mini bead beater (Biospec). After removal of the glass beads, the cell extract was heated in the presence of SDS (1%) at 80°C for 15 minutes. Separately, a negative control was extracted from the transformant carrying pR10M by the same procedure.

[0063] 50 µg protein from the extract was analyzed by SDS-polyacrylamide gel electrophoresis (Figure 11(a) and (b)). After Coomassie Brilliant Blue (CCB) staining, the extract and a recombinant green fluorescent protein (Clontech) as the positive control showed major bands with a molecular weight of 25000, but the negative control did not. Further, 50 µg protein from the extract was analyzed by SDS-polyacrylamide gel electrophoresis followed by western blotting on a PVDF membrane using an anti-green fluorescent protein antibody (Clontech). The cell extract and the positive control showed major bands with a molecular weight of 25000, but the negative control did not. These results provide biological evidence of expression of the desired green fluorescent protein.

EXMAPLE 11

Optimization of the incubation method (1)

[0064] The transformed *S. pombe* strain ASP (Example 9) was incubated in YPD medium containing 100 µg/ml G418 (YPD100) at 32°C. The expression level of the green fluorescent protein in the cell culture was determined fluorometrically by means of a microplate reader (Corona Electric Co., Ltd.) equipped with a fluorescent attachment (excitation wavelength 490nm/ emission wavelength 530nm) (Figure 12(a) and (b)). OD, FLU/OD and time denote the cell density, the fluorescence intensity per cell and the incubation time, respectively. The results show that strain ASP138 did not express the green fluorescent protein variant until the late-growth phase after glucose exhaustion in the mid-growth phase, unlike strain ASP122 having a non-inducible cytomegalovirus promoter [a transformant carrying a recombination product of pHGFPs65T (Clontech) for expression of a green fluorescent protein variant of S65T prepared as disclosed in JP-A-7-163373], clearly due to repression of the inducible invertase promoter in the presence of glucose and subsequent derepression by glucose exhaustion, demonstrating the applicability of this mechanism to expression of the green fluorescent protein as a heterologous protein.

EXAMPLE 12

Incubation of strain ASP138 (2)

5 [0065] The transformed *S. pombe* strain ASP138 (Example 9) was incubated in YPD medium containing 100 µg/ml G418 (glucose concentration 8%) at 32°C to the mid-growth phase, and after medium change, incubated in YPDG medium containing 100 µg/ml G418 (glucose concentration 0.1%, glycerol concentration 3%) (Figure 13(a) and (b)). OD, FLU/OD and time denote the cell density, the fluorescence intensity per cell and the incubation time, respectively. The results show that while the green fluorescent protein was not expressed in the cells incubated without medium change (Untreated), expression of the green fluorescent protein in the cells incubated with the medium change was activated by the medium change (Medium-changed) probably because the depletion of glucose in the medium provoked derepression of the invertase promoter and thereby induced the protein expression. The high level expression induced by the medium change to a low-glucose expression medium suggests that use of a growth medium (with a high glucose concentration) and an expression medium (with a low glucose concentration) can differentiate between cell growth and protein expression. It was demonstrated that the repression of the inducible invertase promoter in the presence of glucose and derepression by exhaustion of glucose could be utilized in expression of the green fluorescent protein as a heterologous protein.

EXAMPLE 13

Incubation of strain ASP138 (3)

20 [0066] The transformed *S. pombe* strain ASP138 (Example 9) was incubated in YPD medium containing 100 µg/ml G418 (glucose concentration 8%) at 32°C to the late-growth phase, and after medium change, incubated in YPDG medium containing 100 µg/ml G418 (glucose concentration 0.1%, glycerol concentration 3%) (Figure 14(a) and (b)). OD, FLU/OD and time denote the cell density, the fluorescence intensity per cell and the incubation time, respectively. The results show that while the green fluorescent protein was not expressed in the cells incubated without medium change (Untreated), expression of the green fluorescent protein in the cells incubated with the medium change was activated by the medium change (Medium-changed) probably because the depletion of glucose in the medium provoked derepression of the invertase promoter and thereby induced the protein expression. The high level expression induced by the medium change to a low-glucose expression medium from a high-glucose medium before glucose exhaustion suggests that use of a growth medium (with a high glucose concentration) and an expression medium (with a low glucose concentration) can differentiate between cell growth and protein expression. It was demonstrated that the repression of the inducible invertase promoter in the presence of glucose and derepression by exhaustion of glucose could be utilized in expression of the green fluorescent protein as a heterologous protein.

EXAMPLE 14

Construction of inducible lipocortin I expression vector pRI0LPI

40 [0067] PCR using plasmid pINV3000 (Example 1) carrying the *S. pombe* invertase gene as the template and oligo DNAs shown in SEQ ID NOS: 14 and 15 in the Sequence Listing as primers gave an amplification product containing the promoter region in the invertase gene and having restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with *Spe*I and *Eco*RI, the amplification product was subjected to agarose gel electrophoresis. Purification of the band of about 3000 bp by the glass beads method using EASY-TRAP gave a fragment for use as a promoter insert.

[0068] The expression vector pTL2L (JP-A-7-163373) carrying a human lipocortin I gene was subjected to agarose gel electrophoresis after terminal double restriction digestion with *Eco*RI and *Hind*III. Purification of the band of about 1000 bp by the glass beads method using EASY-TRAP gave a fragment for use as the OPF insert.

50 [0069] The *S. pombe* multicloning vector pTL2M (JP-A-7-163373) was subjected to agarose gel electrophoresis after terminal double restriction digestion with *Spe*I and *Hind*III. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

[0070] The three fragments were ligated with a DNA ligation kit and transformed into *E. coli* strain DH5. *E. coli* colonies were screened for the inducible lipocortin I expression vector pRI0LPI through base sequencing and restriction mapping, and the vector was recovered by the alkali-SDS method on a preparatory scale.

EXAMPLE 15

Preparation of fission yeast *Schizosaccharomyces pombe* transformant ASP139

[0071] A *S. pombe* wild-type strain ARC001 was transformed with the inducible lipocortin I expression vector pRI0LPI and a transducing vector pAL7 as described by Okazaki et al.

[0072] 1 ml of a preincubated ARC001 culture in YPD medium was incubated in minimum medium MB+Leu with shaking at 30°C for 16 hours to a cell density of 1×10^7 per 1 ml, and the cells were collected, washed with water, suspended in 1 ml of 0.1M lithium acetate (pH 5.0) and incubated at 30°C for 60 minutes. A 100 µl portion of the suspension was mixed with 2 µg of the recombinant vector pRI0LPI and 0.5 µg of PstI-digested pAL7 in 15 µl TE and further with 290 µl of 50% PEG 4000 thoroughly and incubated at 30°C for 60 minutes, at 43°C for 15 minutes and at room temperature for 10 minutes, successively. After centrifugal removal of PEG, the cells were suspended in 1 ml of 1/2 YEL+Leu medium. After 10-fold dilution, 1 ml of the suspension was incubated at 32°C for 2 hours, and a 300 µl portion was spread on minimum medium agar MMA. After 3 days of incubation at 32 °C, about 300 independent colonies had developed on the plate.

[0073] 10 colonies of the transformant were inoculated in 2 ml of YEL medium containing 10 µg/ml antibiotic G418 (YEL10 medium) and incubated with shaking at 32°C. 2 days later, 2 clones were viable. All the subcultures of them were viable 3 days later. The putative desired transformant (ASP138 strain) was frozen in glycerol and stored for use in subsequent experiments.

EXAMPLE 16

Analysis of lipocortin I expression

[0074] *S. pombe* transformant ASP139 (Example 15) was incubated in YPD medium containing 100 µg/ml G418 (glucose concentration 8%) at 32°C to the stationary phase and collected as a non-inducible cell culture. Separately, ASP139 was incubated in the same medium at first to the mid-growth phase, then after medium change, incubated in YPDG medium containing 100 µg/ml G418 (glucose concentration 0.1%, glycerol concentration 3%) to the stationary phase and collected as an inducible cell culture. Both cell cultures were washed, suspended in 50mM tris-HCl (pH 7.5) and disrupted with glass beads in a mini bead beater. After removal of the glass beads, the cell extracts were heated in the presence of SDS (1%) at 80°C for 15 minutes.

[0075] 50 µg protein from each extract was separated by SDS-polyacrylamide gel electrophoresis and stained with Coomassie Brilliant Blue. The extract from the inducible cell culture showed a major band of a molecular weight of about 45000, which is the same as the deduced molecular weight of the recombinant lipocortin I protein, but the extract from the non-inducible cell culture did not. More sensitive western analysis of band density showed that the band from the inducible cell culture extract was 10 times denser than the band from the non-inducible cell culture extract. The results show that while lipocortin I was not expressed in the cells incubated without medium change, the expression of lipocortin I in the cells incubated with the medium change was activated by the medium change probably because the depletion of glucose in the medium provoked derepression of the invertase promoter and thereby induced the protein expression. The high level expression induced by the medium change to a low-glucose expression medium from a high-glucose medium suggests that use of a growth medium (with a high glucose concentration) and an expression medium (with a low glucose concentration) can differentiate between cell growth and protein expression. It was demonstrated that the repression of the inducible invertase promoter in the presence of glucose and derepression by exhaustion of glucose could be utilized in expression of lipocortin I as a heterologous protein.

EXAMPLE 17

Construction of expression vector pTL2INV1 carrying invertase gene

[0076] PCR using plasmid pINV3000 (Example 1) carrying the *S. pombe* invertase gene as the template and oligo DNAs shown in SEQ ID NOS: 16 and 17 in the Sequence Listing as primers gave an amplification product containing the ORF in the invertase gene and having restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with AflIII (New England Biolab) and HindIII (Takara Shuzo Co., Ltd), the amplification product was subjected to agarose gel electrophoresis. Purification of the band of about 3000 bp by the glass beads method using EASY-TRAP gave a fragment for use as an insert.

[0077] The *S. pombe* multicloning vector pTL2M (JP-A-7-163373) was subjected to agarose gel electrophoresis after terminal double restriction digestion with SpeI and EcoI. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

EP 1 223 219 A2

[0078] The two fragments were ligated with a DNA ligation kit (Takara Shuzo Co., Ltd) and transformed into *E. coli* strain DH5 (Toyobo Co., Ltd.). *E. coli* colonies were screened for the invertase gene expression vector pRI0LPI through base sequencing and restriction mapping, and the vector was recovered multiplied by the alkali-SDS method on a preparatory scale.

EXAMPLE 18

Construction of secretory expression vector pSL2I06a'c1 using the signal sequence from the invertase gene

[0079] PCR using plasmid pINV3000 (Example 1) carrying the *S. pombe* invertase gene as the template and oligo DNAs shown in SEQ ID NOS: 18 and 19 in the Sequence Listing as primers gave an amplification product containing the ORF in the invertase gene and having restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with *SpeI* (Takara Shuzo Co., Ltd.) and *EcoRI* (Takara Shuzo Co., Ltd.), the amplification product was subjected to agarose gel electrophoresis. Purification of the band of about 700 bp by the glass beads method using EASY-TRAP (Takara Shuzo Co., Ltd.) gave a fragment for use as a signal insert.

[0080] PCR using plasmid pSL2P06a'c1 (WO96/23890) containing human interleukin 6a'c1 variant cDNA as the template and oligo DNAs shown in SEQ ID NOS: 20 and 21 in the Sequence Listing as primers gave an amplification product containing the interleukin 6a'c1 variant ORF. After terminal double restriction digestion with *EcoRI* and *HindIII* (Takara Shuzo Co., Ltd), the amplification product was subjected to agarose gel electrophoresis. Purification of the band of about 600 bp by the glass beads method using EASY-TRAP (Takara Shuzo Co., Ltd.) gave a fragment for use as a gene insert.

[0081] The *S. pombe* multicloning vector pTL2M (JP-A-7-163373) was subjected to agarose gel electrophoresis after terminal double restriction digestion with *SpeI* and *HindIII*. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

[0082] The three fragments were ligated with a DNA ligation kit and transformed into the *E. coli* strain DH5 (Toyobo Co., Ltd.). *E. coli* colonies were screened for the IL-6a'c1 secretory expression vector pSL2I06a'c1 through base sequencing and restriction mapping, and the vector was recovered by the alkali-SDS method on a preparatory scale.

EXAMPLE 19

Preparation of *S. pombe* transformant ASP168

[0083] A leucine-requiring *S. pombe* strain ARC001 was transformed with the interleukin-6a'c1 variant secretory expression vector pSL2I06a'c1 (Example 18) and a transducing vector pAL7 as described by Okazaki et al.

[0084] 1 ml of a preincubated ARC001 culture in YPD medium was incubated in 100 minimum medium MB+Leu with shaking at 30°C for 16 hours. The cells were collected, washed with water, suspended in 0.1M lithium acetate (pH 5.0) at a cell density of 10^9 cells/ml and incubated at 30°C for 60 minutes. A 100 μ l portion of the suspension was mixed with 2 μ g of the recombinant vector pSL2I06a'c1 and 1.0 μ g of *PstI*-digested pAL7 in 15 μ l TE and further with 290 μ l of 50% PEG 4000 thoroughly and incubated at 30°C for 60 minutes, at 43°C for 15 minutes and at room temperature for 10 minutes, successively. After centrifugal removal of PEG, the cells were suspended in 1 ml of 1/2 YEL+Leu medium. After 10-fold dilution, 1 ml of the suspension was incubated at 32°C for 2 hours, and a 300 μ l portion was spread on minimum medium agar MMA. After 3 days of incubation at 32 °C, about 1000 independent colonies had developed on the plate.

[0085] The transformants (colonies) were inoculated in 2 ml of YEL medium containing 10 μ g/ml antibiotic G418 (YEL10 medium) and incubated with shaking at 32°C for 5 days. The viable clones of the putative desired transformant, designated as strain ASP168, were frozen in glycerol and stored for use in subsequent experiments.

EXAMPLE 20

Analysis of expressed secretory interleukin-6a'c1 variant in culture medium

[0086] A fission yeast *Schizosaccharomyces pombe* transformant ASP168 (Example 19) was incubated in MA-Casamino acid medium (MA medium containing 2% Casamino acid and 3% glucose; for the composition of MA medium, refer to "Alfa et al., Experiments with Fission Yeast, Cold Spring Harbor Laboratory Press, 1993") containing 500 μ g/ml G418 at 32°C for 2 days.

[0087] The cell culture was centrifuged, and the supernatant was concentrated 100-fold through a membrane filter (Amicon Co., Ltd.). Analysis of the concentrated sample by SDS-polyacrylamide electrophoresis followed by Coomassie Brilliant Blue staining gave the SDS-PAGE pattern shown in Figure 15. Lane 1 is the purified interleukin-6a'c1

EP 1 223 219 A2

variant, lane 2 is the supernatant from the ASP168 cell culture, and lane 3 is the supernatant from a cell culture of the control strain ASP021 [transformant carrying a recombinant vector with no ORF prepared by recombination of pTL2M (JP-A-7-163373) by the method disclosed in JP-A-7-163373 without introduction of any gene to be expressed]. The band with a molecular weight of about 20000 in lane 3 seemed attributable to the interleukin-6a'c1 variant from the comparison of lanes 1 and 3.

[0088] Further analysis by western blotting using an anti-IL-6a'c1 gave the pattern shown in Figure 16. Lane 1 is the purified interleukin-6a'c1 variant, lane 2 is the supernatant from the ASP168 cell culture, and lane 3 is the supernatant from a cell culture of the control strain ASP021. The band with a molecular weight of about 20000 in lane 3 was identified as the interleukin-6a'c1 variant from the comparison of lanes 1 and 3.

SEQUENCE LISTING

<110> Asahi Glass Co., Ltd.

<120> Inducible promoter gene and secretion signal gene functional in Schizosaccharomyces pombe, expression vector containing them and their use

<130> AB23(F9865)

<150> JP9/314608

<151> 1997-10-31

<160> 21

<170> MS-DOS

<210> 1

<211> 4748

<212> DNA

<213> Schizosaccharomyces pombe

<400> 1

```

ggatcctagt ccgcgaaatc gagatgcttt gaagattzaa attaaattta attttatgcg      60
agactggttt ctttatTTTT tgtatagtcg catgcaagcg aggttcgcat aatttggaaa      120
ataaaggtag tcaagaagac gttgaattaa ggctgcagtt tcaaagtact ctacaaacga      180
ttccttttta aaaaaaagat tcaaaaaaaa ggcaaagggg ttaagtaatg cttgttattt      240
caatttacct ccaaacagtt actaatgcaa ttgcaaaaaa aaaacctacc tattgaatca      300
aaatttctag cccatccatc gctcctcaag ataaaggaat cgatatTTTg agtttaaggg      360
agttgctgat agatttcaga attaaaaatt ttggaaaaag gatgtcgaga acaagaagat      420
acgtctagat tgctgatgat gcattctagc agacggaaat acaacgatat gtggacagca      480
cgacttttga tccgttcgga tcaaaaggaa gagaaatatt catctttcaa gaagaatgca      540
ggaaaagcaa taaatgcccc ttigtattcct aaattatccc caaaaatgaa cattatgaga      600
tcttcttTgt ggagacagga aatttcgcaa ttccaaacga aaattcggct ctttttttta      660
ccccacagtt gcggggtaaa tgatgtaacg gaccttgggg gaaaggatga tgagttagtt      720
gggaagcgga aaaaatggaa aacggaagta agaatagaaa ccagtatggc tgagtgcatt      780
ggcgggaaaag attttacaga gatgacaaga atctatttat ctataaggaa aaactttttc      840
caaatttTgt taaaaacgca ttctcctcaa ttgcctctag gtagatgata taacgaattg      900
gaacgagaca tcgctaaccg gttttctttg taaatgacat ttgttagtgg gagtaagttt      960
gaatggaggg atagacagat gaatagtatg agatagaaga atagtatata taatgatata      1020
gatgaacaaa taaaaattga aagaaaaaag aaattgttgg ctcatTTgt tcatacacat      1080
gttggttcat acaactttta cccatcgtaa gtattataag taaaaaatag agtacgaaza      1140
gctataagta gtgaagcaaa aaaatagaaa aatagaaaaa aaaatatata taaaaaaata      1200
taataaaaaa aaaactcata agagacgtaa aacacaagaa ttgtctatca ttgtttcttt      1260

```

EP 1 223 219 A2

	aagaagcacc accatctctg aaaactcttc attctctatt agcaaggacc cttttcattc	1320
5	cttctctctt agaatccttt tcattataac gaattggata atcgcgaat aagaacacat	1380
	ccccaaata cgatatatcg atccatcttt tactttgcct agcttattgc tgtacaattc	1440
	catrtaaata gtttctcttc aagaaagatc gicaatggag gcgacaatat accggaattc	1500
10	aagttgcgga cacagagctt gaaaagactg ctttttctat tgttttcaag taaatgaaac	1560
	tgagttttga agtctcaaaa tacatcttat gtattgaaca ttagaagaac atataagata	1620
15	gatcttgaga gctcaattca tcgacattct agccatcata ctgcgatctt agacattgtc	1680
	agcacaacct tagatcgaaa atgaacacgt taccaaacgt tgtctaaaaac ttgccgaatc	1740
	ttatctccgc attacttccg taatccttag tacatacgtt gcaatttcgg aagggtcatga	1800
20	tcgacttttt gtgtagctat aagtgcgca aatgagaaac atgacaagggt gcgatattta	1860
	gcaagatatatt atgcatttga tggagaaaagg aaatttcgga tgtatatata gtaccgttag	1920
	ctgcgctttt ttgggtcctc cataattttc aaactcactg ctttcgatca gatttaccgt	1980
25	ttttaaggtc ttatttgcct tgtgatctgt aggtttggaac atctatagtt cattttctaa	2040
	aagatccttt catcgtttca tcggatagta atcgttcaag aaaaaaaaaag aaaaaaagaa	2100
30	aaagaaaaag aaaagaaaaa taaaccgcta taattcatta cctatttgac tgaaggttct	2160
	tcactttgaa ttgttttgaa tcaaaataaa gaaattatta ttattatttt tttctctgc	2220
	ttttctttta tccattcgtc gaaactattt ttctgtgat aaaagcaatc attccttttt	2280
35	cctgcttctc ttgttatctg aatttttaac gacttttttt cctcgtccat tccctaattc	2340
	tttgcgacct tttctgattc tatccttgggt ttgtactttc gttgtgtaat tgttgagaaa	2400
	gtgaactgat tatttaattg ttgtgaaaaa aattctaaaa ctattttgtt tttcttgatc	2460
40	attcatcctt tgctcgcttg cttgaatatt acagaaattc gtctcccttt caacggaata	2520
	tgataatttg ttgaatactc taaatcaatt aacacctatc aaaagctgaa acattaaatc	2580
45	tattctcacc aaaaaaaaaag actcaagctt cttcgttgtt ggccggtctc ttttttgttt	2640
	tacgattgtt aaattttata ctcaaacctg ccaattctcc acttttgact atttattgat	2700
	agtccttatt taattttctg ttcaccgatt atcgtctttt ttgtaataaa tctttcttgg	2760
50	aaccaaccaa ttaatacgtt ataatcgcta actttgaaga ttgtctaca atg ttt ttg	2818
		Met Phe Leu
		1
55	aaz tat att tta gct agt ggc att tgc ctc gtc tct ctc tta tca tct	2866
	Lys Tyr Ile Leu Ala Ser Gly Ile Cys Leu Val Ser Leu Leu Ser Ser	

EP 1 223 219 A2

	5	10	15	
5	aca aac gcg gct ccc cgt cac tta tat gta aaa cgt tat cct gtc att Thr Asn Ala Ala Pro Arg His Leu Tyr Val Lys Arg Tyr Pro Val Ile 20 25 30 35	2914		
10	tat aat gct tcc aac atc act gaa gtc agc aat tct acc acc gtt cct Tyr Asn Ala Ser Asn Ile Thr Glu Val Ser Asn Ser Thr Thr Val Pro 40 45 50	2962		
15	cct cct cca ttc gta aat aca acg gcc cct aat ggg act tgt ttg ggt Pro Pro Pro Phe Val Asn Thr Thr Ala Pro Asn Gly Thr Cys Leu Gly 55 60 65	3010		
20	aac tat aac gag tat ctt cct tca gga tat tac aat gct acc gat cgt Asn Tyr Asn Glu Tyr Leu Pro Ser Gly Tyr Tyr Asn Ala Thr Asp Arg 70 75 80	3058		
25	ccc aaa att cat ttt act cct tct tcc ggt ttc atg aat gat cca aac Pro Lys Ile His Phe Thr Pro Ser Ser Gly Phe Met Asn Asp Pro Asn 85 90 95	3106		
30	gga ttg gta tat act ggc ggc gtc tat cac atg ttc ttc caa tat tca Gly Leu Val Tyr Thr Gly Gly Val Tyr His Met Phe Phe Gln Tyr Ser 100 105 110 115	3154		
35	cca aaa act cta aca gcc ggc gaa gtt cat tgg ggt cac aca gtt tcc Pro Lys Thr Leu Thr Ala Gly Glu Val His Trp Gly His Thr Val Ser 120 125 130	3202		
40	aag gat tta atc cat tgg gag aat tat cct att gcc atc tac ccc gat Lys Asp Leu Ile His Trp Glu Asn Tyr Pro Ile Ala Ile Tyr Pro Asp 135 140 145	3250		
45	gaa cat gaa aac gga gtt ttg tcc ctc cca ttt agt ggc agt gca gtc Glu His Glu Asn Gly Val Leu Ser Leu Pro Phe Ser Gly Ser Ala Val 150 155 160	3298		
50	gtc gat gtt cat aat tct tcc ggt ctc ttt tcc aac gac acc att ccg Val Asp Val His Asn Ser Ser Gly Leu Phe Ser Asn Asp Thr Ile Pro 165 170 175	3346		
55	gaa gag cgc att gtt tta att tat acc gat cat tgg act ggt gtt gct Glu Glu Arg Ile Val Leu Ile Tyr Thr Asp His Trp Thr Gly Val Ala 180 185 190 195	3394		
	gag cgt cag gct att gcg tat acc act gat ggt gga tat act ttc aaa Glu Arg Gln Ala Ile Ala Tyr Thr Thr Asp Gly Gly Tyr Thr Phe Lys 200 205 210	3442		
	aaa tat tca gga aat ccc gtt ctt gac att aat tca ctt caa ttc cgc Lys Tyr Ser Gly Asn Pro Val Leu Asp Ile Asn Ser Leu Gln Phe Arg 215 220 225	3490		
	gac ccc aag gta ata tgg gat ttc gat gct aat cgt tgg gtg atg att Asp Pro Lys Val Ile Trp Asp Phe Asp Ala Asn Arg Trp Val Met Ile	3538		

EP 1 223 219 A2

	230	235	240	
5	gta gct atg tct caa aat tat gga att gcc ttt tat tcc tcc tat gac Val Ala Met Ser Gln Asn Tyr Gly Ile Ala Phe Tyr Ser Ser Tyr Asp 245 250 255	3586		
10	ttg att cac tgg acc gag tta tct gtt ttc tcc act tct ggt tat ttg Leu Ile His Trp Thr Glu Leu Ser Val Phe Ser Thr Ser Gly Tyr Leu 260 265 270 275	3634		
15	ggg ttg caa tat gaa tgc cct gga atg gct cgt gtg ccc gtt gaa ggc Gly Leu Gln Tyr Glu Cys Pro Gly Met Ala Arg Val Pro Val Glu Gly 280 285 290	3682		
20	acc gat gaa tac aaa tgg gta ctc ttc at ctc catc aat cct ggc gct Thr Asp Glu Tyr Lys Trp Val Leu Phe Ile Ser Ile Asn Pro Gly Ala 295 300 305	3730		
25	cca ttg gga gga tcc gtt gtc caa tac ttt gtt ggc gat tgg aat ggt Pro Leu Gly Gly Ser Val Val Gln Tyr Phe Val Gly Asp Trp Asn Gly 310 315 320	3778		
30	aca aac ttc gtc ccc gat gat ggc caa act aga ttc gta gac ttg ggt Thr Asn Phe Val Pro Asp Asp Gly Gln Thr Arg Phe Val Asp Leu Gly 325 330 335	3826		
35	aag gac ttt tac gcc agc gct ttg tat cac tgc tct tcc gcc aat gcc Lys Asp Phe Tyr Ala Ser Ala Leu Tyr His Ser Ser Ser Ala Asn Ala 340 345 350 355	3874		
40	gat gtt att gga gtt gga tgg gct agc aac tgg caa tac acc aac caa Asp Val Ile Gly Val Gly Trp Ala Ser Asn Trp Gln Tyr Thr Asn Gln 360 365 370	3922		
45	gct cct act caa gtt ttc cgc agt gct atg aca gtt gca cga aaa ttc Ala Pro Thr Gln Val Phe Arg Ser Ala Met Thr Val Ala Arg Lys Phe 375 380 385	3970		
50	act ctt cgc gac gtt cct cag aac ccc atg acc aac ctt act tct ctc Thr Leu Arg Asp Val Pro Gln Asn Pro Met Thr Asn Leu Thr Ser Leu 390 395 400	4018		
55	att caa acc cca ttg aat gtt tct ctc tta cga gat gaa aca cta ttt Ile Gln Thr Pro Leu Asn Val Ser Leu Leu Arg Asp Glu Thr Leu Phe 405 410 415	4066		
60	acc gca ccc gtt atc aat agt tca agt agt ctt tgc ggc tct ccg att Thr Ala Pro Val Ile Asn Ser Ser Ser Ser Leu Ser Gly Ser Pro Ile 420 425 430 435	4114		
65	act ctt cca agc aat acc gca ttc gag ttc aat gtc aca ctc agt atc Thr Leu Pro Ser Asn Thr Ala Phe Glu Phe Asn Val Thr Leu Ser Ile 440 445 450	4162		
70	aat tac aca gaa ggc tgc aca aca gga tat tgt ctg ggg cgt att atc Asn Tyr Thr Glu Gly Cys Thr Thr Gly Tyr Cys Leu Gly Arg Ile Ile	4210		

EP 1 223 219 A2

	455	460	465	
5	att gat tct gat gat cca tac aga tta caa tcc atc tcc gtg gac gtt Ile Asp Ser Asp Asp Pro Tyr Arg Leu Gln Ser Ile Ser Val Asp Val 470 475 480	4258		
10	gat ttt gca gct agc act tta gtc att aat cgt gcc aaa gct cag atg Asp Phe Ala Ala Ser Thr Leu Val Ile Asn Arg Ala Lys Ala Gln Met 485 490 495	4306		
15	gga tgg ttt aat tca ctt ttc acg cct tct ttt gcc aac gat att tac Gly Trp Phe Asn Ser Leu Phe Thr Pro Ser Phe Ala Asn Asp Ile Tyr 500 505 510 515	4354		
20	att tat gga aac gta act ttg tat ggt att gtt gac aat gga ttg ctt Ile Tyr Gly Asn Val Thr Leu Tyr Gly Ile Val Asp Asn Gly Leu Leu 520 525 530	4402		
25	gaa ctg tat gtc aat aat ggc gaa aaa act tac act aat gac ttt ttc Glu Leu Tyr Val Asn Asn Gly Glu Lys Thr Tyr Thr Asn Asp Phe Phe 535 540 545	4450		
30	ttc ctt caa gga gca aca cct gga cag atc agc ttc gct gct ttc caa Phe Leu Gln Gly Ala Thr Pro Gly Gln Ile Ser Phe Ala Ala Phe Gln 550 555 560	4498		
35	ggc gtt tct ttc aat aat gtt acc gtt acg cca tta aag act atc tgg Gly Val Ser Phe Asn Asn Val Thr Val Thr Pro Leu Lys Thr Ile Trp 565 570 575	4546		
40	aat tgc taaatatattt gtttcaagtt aggaaagtat aataactttt gtccttgcac Asn Cys 580	4602		
45	attcaattgt aaagtttagt ttatcctttc atcgtaacca caattgtcac ctaaatctct aaaaatctct tcacttatct agttaatgtc gtaacaaaaa agtccagtag cttcgggaaa tgatgcttgg aatcatacaa gtcgac	4662 4722 4748		
50	<210> 2 <211> 7286 <212> DNA <213> Artificial Sequence <400> 2 agcttgaaaa aacctccac acctccccct gaacctgaaa cataaaatga atgcaattgt tgttgttaac ttgtttattg cagcttataa tggttacaaa taaagcaata gcatcacaaa tttcacaaat aaagcatttt ttctactgca ttctagttgt gggttgtcca aactcatcaa tgtatcttat catgtctgga tcgatcccg gagggtgggc gtcgcttggc cgggtcatttc gaaccccgaga gicccgctca gaagaactcg tcaagaagge gatagaagge gatgcgctgc	60 120 180 240 300		
55				

EP 1 223 219 A2

gaatcggag cggcgatacc gtaaagcag aggaagcgt cagcccatc gccgccaagc 360
5 tcttcagcaa tatcacgggt agccaacgt atgtcctgat agcggtcgc cacacczagc 420
cggccacagt cgalgaatcc agaaaagcgg ccattttcca ccatgatatt cggcaagcag 480
gcatcgccat gggtcacgac gagatcctcg ccgtcgggca tgcgcgcctt gagcctggcg 540
10 aacagttcgg ctggcgcgag cccctgatgc tcttcgtcca gatcatcctg atcgacaaga 600
ccggtttcca tccgagtagc tgcctcgtcg atgcgatgtt tcgcttggtg gtcgaatggg 660
caggtagccg gatcaagcgt atgcagccgc cgcattgcat cagccatgat ggatactttc 720
15 tcggcaggag caaggtgaga tgacaggaga tcttgcctcg gcacttcgcc caatagcagc 780
cagtccttcc ccgttcagt gacaacgtcg agcacagctg cgcaagggaac gcccgtcgtg 840
20 gccagccacg atagccgcgc tgcctcgtcc tgcagttcat tcagggcacc ggacaggtcg 900
gtcttgacaa aaagaaccgg gcgcccctgc gctgacagcc ggaacacggc ggcatcagag 960
cagccgattg tctgttgtgc ccagtcatag ccgaatagcc tctccacca agcggccgga 1020
25 gaacctgcgt gcaatccatc ttgttcaatc atgcgaaacg atcctcatcc tgtctcttga 1080
tcagatccgg gacctgaaat aaaagacaaa aagactaaac ttaccagtta actttctggt 1140
30 ttttcagttc ctcgaggagc tttttgcaaa agcctaggcc tccaaaaaag cctcctcact 1200
acttctggaa tagctcagag gccgaggcgg cctcggcctc tgcataaata aaaaaatta 1260
gtcagccatg gggcggagaa tgggcggaac tgggcggagt tagggcggg atgggcggag 1320
35 ttagggcggg gactatggtt gctgactaat tgagatgcat gctttgcata cttctgcctg 1380
ctggggagcc tggggacttt ccacacctgg ttgctgacta attgagatgc atgctttgca 1440
tacttctgcc tgcctggggag cctggggact tccacaccc taactgacac acattccaca 1500
40 ggacattgat tattgactag ttagtccgcg aaatcgagat gctttgaaga ttaaaattaa 1560
atttaatttt atgcgagact ggtttcccta ttttttgtat agtcgcatgc aagcgaggtt 1620
cgcataattt ggaaaaataa ggtagtcaag aagacgttga attaaggctg cagtttcaaa 1680
45 gtactctaca aacgattcct tttaaaaaaa aagattcaaa aaaaaggcaa agggtttaag 1740
taatgcttgt tatttcaatt tacctccaaa cagttactaa tgcaattgca aaaaaaaaaac 1800
50 ctacctattg aatcaaaatt tctagcccat ccacgcctcc tcaagataaa ggaatcgata 1860
ttttgagttt aaggagttg ctgatagatt tcagaattaa aaatttttgg aaaaggatgt 1920
cgagaacaag aagatacgtc tagattgctg atgatgcatt ctagcagacg gaaatacaac 1980

EP 1 223 219 A2

	gataatgtgga cagcacgact ttgatccgt tcgzaatcaa aggaagagaa atatccatct	2040
	ttcaagaaga atgcaggaaa agcaataaa: gccatttga ttcttaaat atccccaaaa	2100
5	atgaacatta tgagatcttc ttgtgggaga caggaaattt cgcaattcca aacgazaatt	2160
	cggctctttt ttttacccca cagttgcggg gtaaatgatg taacggacct tgggggzaag	2220
10	gatgatgagt tagttgggaa gcggaaaaaa tggaaaacgg aagtaagaat agaaaccagt	2280
	atggctgagt gcaatggcgg aaaagatttt acagagatga caagaatcta tttatctata	2340
	aggaaaaact ttttccaaat ttgtctaaaa acgcattctc ctcaatgcc tctaggtaga	2400
15	tgatataacg aattggaacg agacatcgct aaccggtttt ctttgtaaat gacattttgt	2460
	agtgggagta agtttgaatg gagggataga cagatgaata gtatgagata gaagaatagt	2520
	atatataatg attaagatga acaataaaaa attgaaagaa aaaagaaatt gttggctcat	2580
20	ttggttcata cacatgttgg ttcatataac ttttaccat cgtaagtatt ataagtaaaa	2640
	aatagagtac gaaaagctat aagtagtgaa gcaaaaaaat agaaaaatag aaaaaaaat	2700
25	atatataaaa aaatataata aaaataaaac tcataagaga cgtaaaacac aagaattgtc	2760
	tatcatttgt tctttaagaa gcaccacat tctgtaaaac tcttcatttc tcattagcaa	2820
	ggaccctttt cattccttcc tctttagaat ccttttcatt ataacgaatt ggataatacg	2880
30	caaataagaa cacatccctt aaatacgata tatcgatcca ttttttactt tgcctagctt	2940
	attgctgtac aattccattt aaatagtttc tcttcaagaa agatcgtcaa tggaggcgac	3000
	aatataccgg aatttaagtt gcggacacag agcttgaaaa gactgcattt tgtattgttt	3060
35	tcaagtaaat gaaactgagt ttggaagtct caaaatacat cttatgtatt gaacattaga	3120
	agaacatata agatagatct tgagagctca attcatcgac attctagcca tcatactgcg	3180
40	atcttagaca ttgtcagcac aaccttagat cgaaaatgaa cagttacca aacgttgtct	3240
	aaaacttgcc gaattctatc tccgcattac tccgtaatc cttagtagat acgctgcaat	3300
	ttcggaaggt catgatcgac tttttgtgta gctataagtg acgcaaatga gaaacatgac	3360
45	aagggtcgat atttagcaag atattatgca ttgatggag aaaggaaatt tcggatgtat	3420
	atatagacc gtttagctgcg ctttttttgg tcatacataa ttttcaaact cactgcttcc	3480
	gatcagattt accgttttta aggtctttat tgccttgtga tctgtagggt ggaacatcta	3540
50	tagttcattt tctaaaagat cctttcatcg ttcatcgga tagtaatcgt tcaagaaaaa	3600
	aaaagaaaaa aagaaaaaga aaaagaaaaa aaaaaataaac cgctataatt cattacctat	3660
55		

EP 1 223 219 A2

5 ttgactgaag gtctctcatc ttgaattgtt ttgaatcaaa ataaagaaat tarrattatt 3720
 10 tttttttttt ttgcgttttt ctttatccat tgcgcgaaac tttttttctg ctgataaaag 3730
 15 caatcattcc tttttctctg tttctcttgtt atttcgaattt taaacgactt tttttctctg 3840
 20 tccattccct aattctttgc gaccttttct gattctatcc ttggtttgta ctttcgttgt 3900
 25 gtaattgttg agaaagtga ctgattattt aattgtttgtg aaaaaaatc taaaactatt 3960
 30 ttgtttttct tgatcattca tcttttgctc gcttgcttga atattacaga aattcgtctc 4020
 35 cttttcaacg gaatatgata atttgttgaa tactctaaat caattaacac ctatcaaaag 4080
 40 ctgaaacatt aaatctattc tcaccaaaaa aaaagactca agcttctctg ttgttgcccg 4140
 45 gtctcttttt tgttttacga ttgttaaatt ttatactcac aactgccaat tctccacttt 4200
 50 tgactattta ttgatagtc ctatttaatt ttctgttcac cgattatcgt cttttttgta 4260
 55 aataatcttt cttggaacca accaattaat acgttataat cgctaacttt gaagatttgc 4320
 60 tacaatggca atgggtacag aattcgagct cggtagccgg ggatcctcta gagtcgacct 4380
 65 gcaggcatgc aagcttaaat aggaaagttt ctcaacagg attacagtgt agctacctac 4440
 70 atgctgaaaa atatagcctt taaatcattt ttatattata actctgtata atagagataa 4500
 75 gtccattttt taaaaatgtt tccccaaac cataaaacc tatacaagtt gttctagtaa 4560
 80 caatacatga gaaagatgtc tatgtagctg aaaataaaat gacgtcacia gacaaaaaaa 4620
 85 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaagta 4680
 90 cttcttgagg cggaagaac cagccggatc cagacatgat aagatacatt gatgagtttg 4740
 95 gacaaaccac aactagaatg cagtgaaaaa aatgctttat ttgtgaaatt tgtgatgcta 4800
 100 ttgctttatt tgaaccatt ataagctgca ataaacaagt taacaacaac aattgcattc 4860
 105 attttatgtt tcagggtcag ggggaggtgt gggaggtttt tttaaagcaag taaaacctct 4920
 110 acaaatgttg tatggctgat tatgatccgg ctgcctcgcg cgtttcggtg atgacgggtga 4980
 115 aaacctctga cacatgcagc tcccgagagc ggtcacagct tgtctgtaag cggatgccgg 5040
 120 gagcagacaa gcccgtcagg gcgcgtcagc ggggtgttggc ggggtgtcggg gcgcagccat 5100
 125 gacccagtea cgtagcgata gcggagtgt tacttgctta actatgcggc atcagagcag 5160
 130 attgtactga gagtgcacca tatgcggtgt gaaataccgc acagatgcgt aaggagaaaa 5220
 135 taccgcatca ggcgtcttct cgttctctcg ctactgact cgctgcgctc ggtcgttcgg 5280
 140 ctgcggcgag cggtatcagc tcactcaaaag gcggtaatac ggttatccac agaatacagg 5340

EP 1 223 219 A2

gataacgcag gaaagaatga gcaaaaggcc agcaaaaggc caggaaccgt aaaaaggccg 5400
 5 cgttgcctggc gtttttccat aggctccgcc cccctgacga gcatcacaaa aatcgacgct 5460
 caagtcagag gtggcgaaac ccgacaggac tataaagata ccaggcggtt ccccttgga 5520
 gctccctcgt gcgctctcct gtcccgaccc tgccgcttac cggataacctg tccgcctttc 5580
 10 tcccttcggg aagcgtggcg ctttctcata gctcacgctg taggtatctc agttcgggtg 5640
 aggtcgttcg ctccaagctg ggctgtgtgc acgaaccccc cgttcagccc gaccgctgcg 5700
 ccttatccgg taactatcgt cttgagtcga acccggttaag acacgactta tcgccactgg 5760
 15 cagcagccac tggtaacagg attagcagag cgaggatatgt aggcgggtgct acagagtctt 5820
 tgaagtgggtg gcctaactac ggctacacta gaaggacagt atttgggtatc tgcgctctgc 5880
 tgaagccagt taccttcgga aaaagagttg gtagctcttg atccggcaaa caaaccaccg 5940
 20 ctggtagcgg tggttttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc 6000
 aagaagatcc tttgatcttt tctacggggt ctgacgctca gtggaacgaa aactcacgtt 6060
 25 aagggtttt ggtcatgaga ttatcaaaaa ggaatcttac ctagatcctt ttaaattaaa 6120
 aatgaagttt taaatcaatc taaagtatat atgagtaaac ttgggtctgac agttaccaat 6180
 gcttaatcag tgaggcacct atctcagcga tctgtctatt tcgttcatcc atagtgcct 6240
 30 gactccccgt cgtgtagata actacgatac gggaggggctt accatctggc cccagtgcgtg 6300
 caatgatacc gcgagacca cgctcaccgg ctccagattt atcagcaata aaccagccag 6360
 ccggaagggc cgagcgcaga agtggctctg caactttatc cgcctccatc cagtctatta 6420
 35 attgttgccg ggaagctaga gtaagtagtt cgccagttaa tagtttgccg aacgttggtg 6480
 ccattgctgc aggcatcgtg gtgtcacgct cgtcgtttgg tatggcttca ttcagctccg 6540
 40 gttcccaacg atcaaggcga gttacatgat ccccatgtt gtgcaaaaaa gcggttagct 6600
 ccttcggtcc tccgatcgtt gtcagaagta agttggccgc agtggttatca ctcatggtta 6660
 tggcagcact gcataattct cttactgtca tgccatccgt aagatgcttt tctgtgactg 6720
 45 gtgagtactc aaccaagtca ttctgagaat agtgtatgcg gcgaccgagt tgctcttgcc 6780
 cggcgtcaac acgggataat accgcgccac atagcagaac ttlaaaagtg ctcatcattg 6840
 50 gaaaacgttc ttcggggcga aaactctcaa ggaatctacc gctgttgaga tccagttcga 6900
 tgtaaccac tcgtgcaccc aactgatctt cagcatcttt tactttcacc agcgtttctg 6960
 ggtgagcaaa aacaggaagg caaaatgccg caaaaaaggg aataagggcg acacggaaat 7020

55

EP 1 223 219 A2

gttgaatact cataactcttc ctttttcaat attattgaag catttatcag ggttattgtc 7080
5 tcatgagcgg atacatatct gaatgtatct agaaaaataa acaaataagg gttccgcgca 7140
catttccccg aaaagtgcga cctgacgtct aagaaacat tattatcatg acattaacct 7200
ataaaaaatag gcgtatcacg aggcccttct gtcttcaaga attggtcgac caattctcat 7260
10 gtttgacagc ttatcatcga taagct 7286

<210> 3
15 <211> 27
<212> DNA
<213> Artificial Sequence
<400> 3
gtttgaattc atgaaygayg gnaaygg 27

20 <210> 4
<211> 24
<212> DNA
<213> Artificial Sequence
25 <400> 4
gtttgaattc yrytggggnc aygc 24

30 <210> 5
<211> 27
<212> DNA
<213> Artificial Sequence
<400> 5
gtttgaattc acyttnnggrt cncgraa 27

35 <210> 6
<211> 24
<212> DNA
<213> Artificial Sequence
40 <400> 6
gatgggtaaa actagtatga acca 24

45 <210> 7
<211> 23
<212> DNA
<213> Artificial Sequence
<400> 7
ttgcgaatta ctagtctccc aca 23

50 <210> 8
<211> 29
<212> DNA
<213> Artificial Sequence
55 <400> 8

EP 1 223 219 A2

gggactagtt agtccgcgaa atcgagatg 29

5 <210> 9
<211> 42
<212> DNA
<213> Artificial Sequence
<400> 9

10 gaggaattct gataccattg ccattgtagc aaatcttcaa ag 42

<210> 10
<211> 30
<212> DNA
15 <213> Artificial Sequence
<400> 10
aaagctagcc attgtagcaa atcttcaaag 30

<210> 11
<211> 27
<212> DNA
20 <213> Artificial Sequence
<400> 11
25 atggctagca agggcgagga gctgttc 27

<210> 12
<211> 27
30 <212> DNA
<213> Artificial Sequence
<400> 12
cccaagctta cttgtacagc tcgtcca 27

<210> 13
<211> 7938
35 <212> DNA
<213> Artificial Sequence
<400> 13

40 agcttgaaaa aacctccac acctccccc gaacctgaaa cataaaatga atgcaattgt 60
tgttgttaac ttgtttattg cagcttataa tgggtacaaa taaagcaata gcatcacaaa 120
tttcacaaat aaagcatttt tttcactgca ttctagtgtt gggttgtcca aactcatcaa 180

45 tgtatcttat catgtctgga tcatcccg gagggtgggc gtcgcttggc cggctatttc 240
gaaccccaga gtcccgctca gaagaactcg tcaagaaggc gatagaaggc gatgcgctgc 300
gaatcgggag cggcgatacc gtaaagcacg aggaagcggc cagcccatte gccgccaagc 360

50 tcttcagcaa tatcaggggt agccaacgct atgtcctgat agcggtcgc caccaccagc 420
cggccacagt cgatgaatcc agaaaagcgg ccattttcca ccatgatatt cggcaagcag 480

55

EP 1 223 219 A2

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

gcatcgccat gggtcacgac gagatccctcg ccgtcgggca tgcgcgcctt gagcctggcg 540
 aacagttcgg ctggcgcgag cccctgatgc tcttcgtcca gatcatcctg atcgacaaga 600
 ccggcttcca tccgagtagc tgcctgcctcg atgcgatgtt tcgcttgggtg gtctaatggg 660
 caggtagccg gatcaagcgt atgcagccgc cgcattgcat cagccatgat ggatactttc 720
 tcggcaggag caaggtgaga tgacaggaga tccctccccg gcacttcgcc caatagcagc 780
 cagtcctttc ccgcttcagt gacaacgtcg agcacagctg cgcaagggaac gcccgtcgtg 840
 gccagccacg atagccgcgc tgcctcgtcc tgcagttcat tcagggcacc ggacaggtcg 900
 gtcttgacaa aaagaaccgg gcgcccctgc gctgacagcc ggaacacggc ggcatcagag 960
 cagccgattg tctgttgtgc ccagtcatag ccgaatagcc tctccacca agcggccgga 1020
 gaacctgcgt gcaatccatc ttgttcaatc atgcgaaacg atcctcatcc tgtctcttga 1080
 tcagatccgg gacctgaaat aaaagacaaa aagactaaac ttaccagtta actttctggg 1140
 ttttcagttc ctgaggagc tttttgcaaa agcctaggcc tccaaaaaag cctctcact 1200
 actttcggaa tagctcagag gccgaggcgg cctcggcctc tgcataaata aaaaaaatta 1260
 gtcagccatg gggcggagaa tgggcggaac tgggcggagt tagggcgggg atgggcggag 1320
 ttagggcgcg gactatgggt gctgactaat tgagatgcat gctttgcata cttctgcctg 1380
 ctggggagcc tggggacttt ccacacctgg ttgctgacta attgagatgc atgctttgca 1440
 tacttctgcc tgcitggggag cctggggact ttccacaccc taactgacac acattccaca 1500
 ggacattgat tatigactag ttagtccgcg aaatcgagat gctttgaaga ttaaaattaa 1560
 atttaatttt atgcgagact ggtttcctta tttttgtat agtcgcatgc aagcgaggtt 1620
 cgcataattt ggaaaataaa ggtagtcaag aagacgttga attaaggctg cagtttcaaa 1680
 gtactctaca aacgattcct tttaaaaaaa aagattcaaa aaaaaggcaa agggtttaag 1740
 taalgtttgt tatttcaatt tacctccaaa cagttactaa tgcaattgca aaaaaaaaaac 1800
 ctacctattg aatcaaaatt tctagcccat ccacgcctcc tcaagataaa ggaatcgata 1860
 ttttgagttt aaggaggttg ctgatagatt tcagaattaa aaatttttgg aaaaggatgt 1920
 cgagaacaag aagatacgtc tagattgctg atgatgcatt ctacgagacg gaaatacaac 1980
 gatatttgga cagcacgact ttgatccgt tcggatcaaa aggaagagaa atatccatct 2040
 ttcaagaaga atgcaggaaa agcaataaat gcccatttga ttccataaatt atccccaaaa 2100
 atgaacatta tgagatcttc ttgtgggaga caggaaattt cgcaattcca aacgaaaatt 2160

EP 1 223 219 A2

5 cggctctttt ttttacccca cagttgcggg gtaaatgatg taacggacct tgggggaaaag 2220
 gatgatgagt tagttgggaa gcggaaaaaa tggaaaacgg aagtaagaat agaaaccagt 2280
 atggctgagt gcaatggcgg aaaagatttt acagagatga caagaatcta tttatctata 2340
 aggaaaaact ttttccaaat ttgtctaaaa acgcattctc ctcaattgcc tctaggtaga 2400
 10 tgatataacg aattggaacg agacatcgct aaccggtttt ctttgtaaat gacattttgt 2460
 agtgggagta agtttgaatg gagggataga cagatgaata gtatgagata gaagaatagt 2520
 atatataatg attaagatga acaataaaaa attgaaagaa aaaagaaatt gttggctcat 2580
 15 ttggttcata cacatgttgg ttcatacaac ttttaccat cgtaagtatt ataagtaaaa 2640
 aatagagtac gaaaagctat aagtagtgaa gcaaaaaaat agaaaaatag aaaaaaaat 2700
 20 atatataaaa aaatataata aaaataaaac tcataagaga cgtaaaacac aagaattgtc 2760
 tatcatttgt tctttaagaa gcaccaccat tctgtaaaac tcttcatttc tcattagcaa 2820
 ggaccctttt cattccttcc tctttagaat ctttttcatt ataacgaatt ggataatacg 2880
 25 caaataagaa cacatccctt aaatacgata tatcgatcca ttttttactt tgcctagctt 2940
 attgctgtac aattccattt aaatagtttc tcttcaagaa agatcgtaa tggaggcgac 3000
 30 aatataccgg aatttaagtt gcggacacag agcttgaaaa gactgcattt tgtattgttt 3060
 tcaagtaaat gaaactgagt tttgaagctt caaaatacat cttatgtatt gaacattaga 3120
 agaacatata agatagatct tgagagctca attcatcgac attctagcca tcatactgcg 3180
 35 atcttagaca ttgtcagcac aaccttagat cgaaaatgaa cacgttacca aacgttgtct 3240
 aaaacttgcc gaatcttate tccgcaattac tccgtaatc cttagtacat acgctgcaat 3300
 ttcggaaggt catgatcgac tttttgtgta gctataagtg acgcaaatga gaaacatgac 3360
 40 aagggtcgat atttagcaag atattatgca ttgtatggag aaaggaaatt tcggatgcat 3420
 atatagtacc gtttagctcg ctttttttgg tcatccataa ttttcaaact cactgctttc 3480
 gatcagattt accgttttta aggtctttat tgctttgtga tctgtagggt ggaacatcta 3540
 45 tagttcattt tctaaaagat cttttcatcg tttcatcgga tagtaatcgt tcaagaaaaa 3600
 aaaagaaaaa aagaaaaaga aaaagaaaag aaaaaataac cgctataatt cattacctat 3660
 50 ttgactgaag gttcttcatc ttgaattgtt tgaatcaaa ataaagaaat tattattatt 3720
 attttttttc ttcgcttttt ctttatccat tcgtcgaaac tatttttctg ctgataaaaag 3780
 caatcattcc ttttctctgc tctctttgtt attcgaaatt taaacgactt ttttctctcg 3840

55

EP 1 223 219 A2

tccattccct aattctttgc gaccttttct gattctatcc ttggtttga ctttcgttgt 3900
 gtaattgttg agaaagtga ctagattatt aattgttttg aaaaaaatc taaaactatt 3960
 5 ttgtttttct tgatcatca tctttgtct gcttgcttga atattacaga aattcgtctc 4020
 cctttcaacg gaatatgata atttgttgaa tactctaaat caattaacac ctatcaaaag 4080
 10 ctgaaacatt aaatctattc tcaccaaaaa aaaagactca agcttcttcg ttgttggccg 4140
 gtctcttttt tgttttacga ttgttaaatt ttatactcac aactgccaat tctccacttt 4200
 tgactattta ttgatagtcc ctatttaatt ttctgttcac cgattatcgt ctttttttga 4260
 15 aataatcttt ctltggaacca accaattaat acgttataat cgctaacttt gaagatttgc 4320
 tacaatggct agcaagggcg aggagctgtt caccgggggtg gtgcccattc tggtcgagct 4380
 ggacggcgac gtaaacggcc acaagttcag cgtgtccggc gagggcgagg gcgatgccac 4440
 20 ctacggcaag ctgacctga agttcatctg caccaccggc aagctgcccg tgccttgccc 4500
 caccctcgtg accaccctga cctacggcgt gcagtgtctc agccgctacc ccgaccacat 4560
 25 gaagcagcac gactttctca agtcgcccat gcccgaggc tacgtccagg agcgcaccat 4620
 cttcttcaag gacgacggca actacaagac ccgcgccgag gtgaagtctg agggcgacac 4680
 cctggtgaac cgcacgagc tgaagggcat cgacttcaag gaggacggca acatcctggg 4740
 30 gcacaagctg gagtacaact acaacagcca caacgtctat atcatggccg acaagcagaa 4800
 gaacggcatc aaggtgaact tcaagatccg ccacaacatc gaggacggca gcgtgcagct 4860
 cgccgaccac taccagcaga acacccccat cggcgacggc cccgtgctgc tgcctgacaa 4920
 35 ccactacctg agcaccagct ccgccctgag caaagacccc aacgagaagc gcgatcacat 4980
 ggtcctgctg gagttcgtga ccgccgccgg gatcactctc ggcatggacg agctgtacaa 5040
 40 gtaagcttaa ataggaaggt tctttcaaca ggattacagt gtagctacct acatgctgaa 5100
 aaatatagcc tttaaatcat ttttatatta taactctgta taatagagat aagtccattt 5160
 tttaaaaatg ttttcccaa accataaaac cctatacaag ttgttctagt aacaatacat 5220
 45 gagaaagatg tctatgtagc tgaaaataaa atgacgtcac aagacaaaaa aaaaaaaaaa 5280
 aaaaaaaaaa aaaaaaaaaa aaaaaaaa aaaaaaaaaa aaaaaaaaag taccttctga 5340
 50 ggcggaaga accagccgga tccagacatg ataagatata ttgatgagtt tggacaaacc 5400
 acaactagaa tgcagtga aaatgtctt atttgtgaa ttgtgatgc tattgcttta 5460
 ttgttaacca ttataagctg caataaaca gttacaaca acaattgcat tcattttatg 5520

55

EP 1 223 219 A2

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

tttcagggttc aggggggaggt gtgggaggtt ttttaaagca agtaaaaccc ctacaaatgt 5580
 ggtagggctg attatgatcc ggctgcctcg cgcgtttcgg tgatgacggg gaaaacctct 5640
 gacacatgca gctcccggag acggtcacag cttgtctgta agcggatgcc gggagcagac 5700
 aagcccgtca gggcgcgtca gcgggtgttg gcgggtgtcg gggcgcagcc atgaccaggt 5760
 cacgtagcga tagcggagtg tatactggct taactatgcg gcatcagagc agattgtact 5820
 gagagtgcac catatgcggt gtgaaatacc gcacagatgc gtaaggagaa aataccgcat 5880
 caggcgcctct tccgcttccct cgctcactga ctgcctgcgc tcggctcgttc ggctgcggcg 5940
 agcggtatca gctcactcaa aggcggtaat acggttatcc acagaatcag gggataacgc 6000
 aggaaagaat gagcaaaaagg ccagcaaaaag gccaggaacc gtaaaaaggc cgcgttgtctg 6060
 gcgtttttcc ataggctccg ccccccctgac gagcatcaca aaaatcgacg ctcaagtcag 6120
 aggtggcgaa acccgacagg actataaaga taccaggcgt tccccctgg aagctccctc 6180
 gtgcgctctc ctgttccgac cctgcgcctt accggatacc tgtccgcctt tctcccttcg 6240
 ggaagcgtgg cgctttctca tagctcacgc ttaggtatc tcagttcggg ttaggtcgtt 6300
 cgctccaagc tgggctgtgt gcacgaacc cccgttcagc ccgaccgtg cgccttatcc 6360
 ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact ggcagcagcc 6420
 actggtaaca ggattagcag agcggaggtat gtaggcggtg ctacagagtt cttgaagtgg 6480
 tggcctaact acggctacac tagaaggaca gtaattggta tctgcgctct gctgaagcca 6540
 gttacccttcg gaaaaagagt tggtagctct tgatccggca aacaaaccac cgctggtagc 6600
 ggtgggtttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc tcaagaagat 6660
 cctttgatct tttctacggg gtctgacgct cagtggaaacg aaaactcacg ttaagggatt 6720
 ttggatcatga gattatcaaa aaggatcttc acctagatcc ttttaaatta aaaatgaagt 6780
 tttaaatcaa tctaaagtat atatgagtaa acttgggtctg acagttacca atgcttaatc 6840
 agtgaggcac ctatctcagc gatctgtcta tttcgctcat ccatagttgc ctgactcccc 6900
 gtcgttaga taactacgat acgggagggc ttaccatctg gccccagtgc tgcaatgata 6960
 ccgcgagacc cacgctcacc ggctccagat ttatcagcaa taaaccagcc agccggaagg 7020
 gccgagcgca gaagtgggtc tgcaacttta tccgcctcca tccagttctat taattgttgc 7080
 cgggaagcta gagtaagtag ttccgaggtt aatagtttgc gcaacgttgt tgccattgct 7140
 gcaggcatcg tgggtgtcag ctgcgtcgtt ggtatggctt cattcagctc cggttcccaa 7200

EP 1 223 219 A2

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

cgaacaaaggc gagttacatg atcccccatg ttgtgcaaaa aagcggttag ctccctcggg 7260
 cctccgatcg ttgtcagaag taagtggcc gcagtgttat cactcatggg tatggcagca 7320
 ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac tgggtgagtac 7380
 tcaaccaagt cattctgaga atagtgtatg cggcgaccga gttgctcttg cccggcgctca 7440
 acacgggata ataccgcgcc acatagcaga actttaaaag tgctcatcat tggaaaacgt 7500
 tcttcggggc gaaaactctc aaggatctta ccgctgttga gatccagttc gatgtzaacc 7560
 actcgtgcac ccaactgac ttcagcatct tttactttca ccagcgttc tgggtgagca 7620
 aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggc cgacacggaa atgttgaata 7680
 ctcatactct tcctttttca atattattga agcatttatt agggttattg tctcatgagc 7740
 ggatacatat ttgaatgtat ttagaaaaat aaacaaatag gggttccgcg cacatttccc 7800
 cgaaaagtgc cacctgacgt ctaagaaacc attattatca tgacattaac ctataaaaaat 7860
 aggcgtatca cgaggccctt tcgtcttcaa gaattggctg accaattctc atgtttgaca 7920
 gcttatcacc gataagct 7938
 <210> 14
 <211> 29
 <212> DNA
 <213> Artificial Sequence
 <400> 14
 gggactagtt agtccgcgaa atcgagatg 29
 <210> 15
 <211> 42
 <212> DNA
 <213> Artificial Sequence
 <400> 15
 gaggaattct gataccattg ccattgtagc aaatcttcaa ag 42
 <210> 16
 <211> 28
 <212> DNA
 <213> Artificial Sequence
 <400> 16
 gggac atgtt tttga aatat atttt agc 28
 <210> 17
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <400> 17

gggaa gctta gcaat tccag atagt ct

27

5

<210> 18
 <211> 20
 <212> DNA
 <213> Artificial Sequence
 <400> 18
 ttgac tagtt attaa tagta

20

10

15

<210> 19
 <211> 29
 <212> DNA
 <213> Artificial Sequence
 <400> 19
 cccga attca taacg tttta catat aagt

29

20

25

<210> 20
 <211> 29
 <212> DNA
 <213> Artificial Sequence
 <400> 20
 aaaga attcc cagta ccccc aacct cttca

29

30

35

<210> 21
 <211> 20
 <212> DNA
 <213> Artificial Sequence
 <400> 21
 aaaat gattt aaagg ctata

20

40

Claims

1. A peptide having the amino acid sequence of Met Phe Leu Lys Tyr Ile Leu Ala Ser Gly Ile Cys Leu Val Ser Leu Leu Ser Ser Thr Asn Ala.

45

2. A DNA encoding the peptide of claim 1.

3. A recombinant vector containing the sequence of the DNA according to claim 2.

50

4. A multicloning vector containing the sequence of the DNA according to claim 2 and a multicloning site.

5. An expression vector containing the sequence of the DNA according to claim 2 and a heterologous protein structural gene.

55

6. A *Schizosaccharomyces pombe* transformant carrying the expression vector according to claim 5.

7. A process for producing a protein which comprises incubating the transformant according to claim 6 and recovering an expressed heterologous protein.

F I G . 1

```

S. pombe inv1      31 : TTAPNGTCLGNYNEYLPSCGYINATDRPKIHFTPSSSGFMNDPNGLVY--TG
S. occidentalis   11 : PLTTTFGGIVASSSIDLSVDTSEYNRPLIHFTPEKGMNDPNGTFYDKTA
S. cerevisiae SUC2  3 : LQAFFTLAGFAAKHSASHTNETSDRPLVHFTPNKGWMNDPGLWIDEKD
                      ..  ....  *  ....  .

S. pombe inv1      101 : GVYHMFQYSPKILTAGE-VHGHGTVSKDLIHWENYPIAIYPDEHENGVL
S. occidentalis    61 : KTWHLFYQYNPNATAHGQPLYWGHATSNDLVHWDHEMAIGPEHNEGIF
S. cerevisiae SUC2  53 : AKWHTYFQYNPNQTVWGTPLFWGHATSDDLTNWEOQPIAIAPKRNDSC--
                      *  ..  .  *  ....  *  ..  .

S. pombe inv1      151 : SLPFSGSIVVDVHN-----SSGLFSNDTI-PEERIVLIYTDHWTGVAERO
S. occidentalis    111 : ----SGSIVVDHNN-----TSGFF-NSSIDPNQRIVAIYTNHMPDL-QTO
S. cerevisiae SUC2  103 : --AFSGSHVVDYNN-----TSGFF-NDTIDPRQRCVAIWT-YNTPESSEEQ
                      ....  *  ..  .  *  ..  .  *  ..  .

S. pombe inv1      201 : AIAITTDGGYTFKKYSGNPVLDINSLQFRDPKVMHDFDANRWVIVAMSQ
S. occidentalis    161 : DIAFSLDCCYTFTRYENNPVIDVSSNQFRDPKVFHHERFKSHDHGCSEIA
S. cerevisiae SUC2  153 : YISYSTDGGYTFTEYQRNPVLAANSTQFRDPKVFHYEP6QRWIMTAARSQ
                      *  ....  *  ..  .  ....  .


S. pombe inv1      251 : NYGIAFYSSYDLIHHTELS VFSTSGYLGLQYECPCGHARVPVEGTDEYK-W
S. occidentalis    211 : RVKIQIFG3ANLKNHVLNSNPS-SGYGNQYGM3RLIEVPIENS3DKSK-W
S. cerevisiae SUC2  203 : DYKIEIYSSDDLKSHKTESAFANECFLCYQYECPCGLIEVPTEQDPSKBYH
                      *  *  *  *  *  *  *  *  *  *  *  *  *  *  *

S. pombe inv1      301 : VLFISINPGAP-LGGSVVQYFVGDNWGTNFPVDDGQTRFVDLGKDFYA-S
S. occidentalis    261 : VMFLAINPGSP-LGGSINQYFVGDFDGFQFVPDD3QTRFVDIGKDFYA-F
S. cerevisiae SUC2  253 : VMFISINPGAP-AGGSFNQYFVG3FNQTHF3AF3DNQ3RVVDFGKDIYALQ
                      *  ....  *  *  *  *  *  *  *  *  *  *  *  *

S. pombe inv1      351 : ALIHSSSANADVIGVGHASHHXYTNQAPTQVF-R3AMTVARKFTLRDVPQ
S. occidentalis    311 : QTF--SEVEHGVGLAWASHHXYADQVPTNPH-R3STSLARNYTLRYVHQ
S. cerevisiae SUC2  303 : TFFNTDPTYGSALGIAWASHHXY3AFVPTNPH-R3BMSLVRKFSLNT3YQ
                      ....  *  *  *  *  *  *  *  *  *  *  *  *

```

FIG. 2



F I G . 3




FIG. 4





FIG. 5

(a)



(b)





FIG. 6



F I G . 7




FIG. 8

invertase activity (U/OD)

repression depression

wild type

$\Delta inv1+$

1.0 kb

$\Delta (-2810/-2810)$

$\Delta (-2810/-2190)$

$\Delta (-2810/-1710)$

$\Delta (-2810/-1200)$

$\Delta (-2810/-200)$

-2000 -1000 0 1000

FIG. 9




FIG. 10




FIG. 11





FIG. 12

(a)



(b)




FIG. 13





FIG. 14





FIG. 15






FIG. 16





Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) EP 1 223 219 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
23.10.2002 Bulletin 2002/43

(43) Date of publication A2:
17.07.2002 Bulletin 2002/29

(21) Application number: 02004252.9

(22) Date of filing: 30.10.1998

(51) Int Cl.7: C12N 15/31, C07K 14/39,
C12N 15/81, C12N 1/19,
C12P 21/02
// (C12N15/31, C12R1:645),
(C12N1/19, C12R1:645)

(84) Designated Contracting States:
DE FR GB

(30) Priority: 31.10.1997 JP 31460897

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
98950470.9 / 0 972 834

(71) Applicant: ASAHI GLASS COMPANY LTD.
Tokyo 100-8405 (JP)

(72) Inventors:

- Tohda, Hideki, Asahi Glass Company Ltd.
Yokohama-shi, Kanagawa 221-0863 (JP)
- Hama, Yuko, Asahi Glass Company Ltd.
Yokohama-shi, Kanagawa 221-0863 (JP)
- Kumagai, Hiromichi, Asahi Glass Company Ltd.
Yokohama-shi, Kanagawa 221-0863 (JP)

(74) Representative: Müller-Boré & Partner
Patentanwälte
Grafinger Strasse 2
81671 München (DE)

(54) Inducible promoter and secretion signal for use in Schizosaccharomyces pombe, expression vector containing them and their use

(57) The present invention relates to an inducible promoter gene for use in the fission yeast Schizosaccharomyces pombe, an inducible expression vector containing the inducible promoter gene, Schizosaccharomyces pombe transformed with the inducible expression vector and a process for producing a heterologous protein by using the Schizosaccharomyces pombe transformant.

The present invention also relates to a secretion signal gene for use in Schizosaccharomyces pombe, an expression vector containing the secretion signal, Schizosaccharomyces pombe transformed with the expression vector and a process for producing a heterologous protein by using the Schizosaccharomyces pombe transformant. These inducible promoter gene and secretion signal gene are originated from the S. pombe gene involved in expression of the invertase precursor.

The inducible promoter gene is originated from the region responsible for the catabolite repression of the

invertase precursor. When a Schizosaccharomyces pombe transformant carrying the inducible expression vector is incubated, the timing of the production of the desired heterologous protein can be controlled by regulating the expression of the heterologous protein structural gene depending on the presence or absence of a specific nutrient (especially glucose).

The secretion signal gene encodes the secretion signal for the invertase precursor. When a Schizosaccharomyces pombe transformant carrying an expression vector containing the secretion signal gene is incubated, a heterologous protein fused with the secretion signal is synthesized as a precursory protein in the cell, and after intracellular separation of the secretion signal, the heterologous protein is secreted from the cell. Therefore, it is possible to obtain the desired protein from the cell culture, and the desired heterologous protein can be purified easily.

EP 1 223 219 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 00 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	MORENO S ET AL: "PURIFICATION AND CHARACTERIZATION OF THE INVERTASE FROM SCHIZOSACCHAROMYCES POMBE" BIOCHEMICAL JOURNAL, PORTLAND PRESS, LONDON, GB, vol. 267, 1990, pages 697-702, XP002917397 ISSN: 0264-6021 * the whole document *	1-7	C12N15/31 C07K14/39 C12N15/81 C12N1/19 C12P21/02 //(C12N15/31, C12R1:645), (C12N1/19, C12R1:645)
X	DATABASE EMBL 'Online! 3 January 1997 (1997-01-03) YOSHIOKA, S.: "Schizosaccharomyces pombe mRNA, partial cds, clone: SY 1543" retrieved from EBI Database accession no. SPD242 XP002169853 * abstract *	1-7	
P, X	TANAKA N ET AL: "ISOLATION AND CHARACTERIZATION OF AN INVERTASE AND ITS REPRESSOR GENES FROM SCHIZOSACCHAROMYCES POMBE" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 245, no. 1, April 1998 (1998-04), pages 246-253, XP002917398 ISSN: 0006-291X * page 246, right-hand column, last paragraph * * figure 1 *	1-7	TECHNICAL FIELDS SEARCHED (Int.Cl.7) C12N C07K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 August 2002	Examiner Schmitz, T
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 92 (P04C011)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 00 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.CI.7)
A	<p>NISHIZAWA M ET AL: "CONSTRUCTION OF AN EXPRESSION AND SECRETION VECTOR FOR THE YEAST SACCHAROMYCES-CEREVISIAE" AGRICULTURAL AND BIOLOGICAL CHEMISTRY, vol. 51, no. 2, 1987, pages 515-521, XP002168488 ISSN: 0002-1369 * page 517, right-hand column, last paragraph - page 518, left-hand column, paragraph 2 * * figure 1 * * page 517, left-hand column, paragraph 3 - right-hand column, paragraph 2 *</p>		
A	<p>PEREZ J A ET AL: "CLONING AND SEQUENCE ANALYSIS OF THE INVERTASE GENE INV1 FROM THE YEAST PICHIA ANOMALA" CURRENT GENETICS, NEW YORK, NY, US, vol. 29, 1996, pages 234-240, XP002917395 ISSN: 0172-8083 * the whole document *</p>		<p>TECHNICAL FIELDS SEARCHED (Int.CI.7)</p>
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 August 2002	Examiner Schmitz, T
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 92 (F04C01)

THIS PAGE BLANK (USPTO)